



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 7 : C12N 15/86, 5/10, A61K 48/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/15820 (43) International Publication Date: 23 March 2000 (23.03.00)</p>
<p>(21) International Application Number: PCT/US99/20118 (22) International Filing Date: 10 September 1999 (10.09.99) (30) Priority Data: 60/099,791 10 September 1998 (10.09.98) US Not furnished 9 September 1999 (09.09.99) US (71) Applicant (for all designated States except US): CALYDON, INC. [US/US]; 1324 Chesapeake Terrace, Sunnyvale, CA 94089 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): YU, De, Chao [CN/US]; 1046 Eagle Lane, Foster City, CA 94404 (US), HENDERSON, Daniel, R. [US/US]; 955 Matadero Avenue, Palo Alto, CA 94306 (US). (74) Agents: POLIZZI, Catherine, M. et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: ADENOVIRUS VECTORS CONTAINING CELL STATUS-SPECIFIC RESPONSE ELEMENTS AND METHODS OF USE THEREOF</p> <div data-bbox="205 730 830 789"> </div> <p>(57) Abstract</p> <p>The present invention provides adenoviral vectors comprising cell status-specific transcriptional regulatory elements which confer cell status-specific transcriptional regulation on an adenoviral gene. A "cell status" is generally a reversible physiological and/or environmental state. The invention further provides compositions and host cells comprising the vectors, as well as methods of using the vectors.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**ADENOVIRUS VECTORS CONTAINING CELL STATUS-SPECIFIC RESPONSE
ELEMENTS AND METHODS OF USE THEREOF**

5 **CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims the priority benefit of U.S. Provisional Patent Application No. 60/099,791, filed September 10, 1998. The priority application is hereby incorporated herein by reference in its entirety.

10 **STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER**

FEDERALLY SPONSORED RESEARCH

(Not Applicable)

TECHNICAL FIELD

15 This invention relates to cell transfection using adenoviral vectors. More specifically, it relates to cell status-specific replication of adenovirus vectors in cells, regardless of tissue or cell type.

BACKGROUND ART

20 In spite of numerous advances in medical research, cancer remains the second leading cause of death in the United States. In the industrialized nations, roughly one in five persons will die of cancer. Traditional modes of clinical care, such as surgical resection, radiotherapy and chemotherapy, have a significant failure rate, especially for solid tumors. Neoplasia resulting in benign tumors can usually be completely cured by removing the mass surgically.

25 If a tumor becomes malignant, as manifested by invasion of surrounding tissue, it becomes much more difficult to eradicate. Once a malignant tumor metastasizes, it is much less likely to be eradicated.

Excluding basal cell carcinoma, there are over one million new cases of cancer per year in the United States alone, and cancer accounts for over one half million deaths per year in this

country. In the world as a whole, the five most common cancers are those of lung, stomach, breast, colon/rectum, and uterine cervix, and the total number of new cases per year is over 6 million.

5 Lung cancer is one of the most refractory of solid tumors because inoperable cases are up to 60% and the 5-year survival is only 13%. In particular, adenocarcinomas, which comprise about one-half of the total lung cancer cases, are mostly chemo-radioresistant. Colorectal cancer is the third most common cancer and the second leading cause of cancer deaths. Pancreatic cancer is virtually always fatal. Thus, current treatment prospects for many patients with these carcinomas are unsatisfactory, and the prognosis is poor.

10 Hepatocellular carcinoma (HCC or malignant hepatoma) is one of the most common cancers in the world, and is especially problematic in Asia. Treatment prospects for patients with hepatocellular carcinoma are dim. Even with improvements in therapy and availability of liver transplant, only a minority of patients are cured by removal of the tumor either by resection or transplantation. For the majority of patients, the current treatments remain
15 unsatisfactory, and the prognosis is poor.

Breast cancer is one of the most common cancers in the United States, with an annual incidence of about 182,000 new cases and nearly 50,000 deaths. In the industrial nations, approximately one in eight women can expect to develop breast cancer. The mortality rate for breast cancer has remained unchanged since 1930. It has increased an average of 0.2% per year, but decreased in women under 65 years of age by an average of 0.3% per year. See e.g., Marchant (1994) *Contemporary Management of Breast Disease II: Breast Cancer*, in: *Obstetrics and Gynecology Clinics of North America* 21:555-560; and Colditz (1993) *Cancer Suppl.* 71:1480-1489.

25 Despite ongoing improvement in the understanding of the disease, breast cancer has remained resistant to medical intervention. Most clinical initiatives are focused on early diagnosis, followed by conventional forms of intervention, particularly surgery and chemotherapy. Such interventions are of limited success, particularly in patients where the tumor has undergone metastasis. There is a pressing need to improve the arsenal of therapies available to provide more precise and more effective treatment in a less invasive way.

Prostate cancer is the fastest growing neoplasm in men with an estimated 244,000 new cases in the United States being diagnosed in 1995, of which approximately 44,000 deaths will result. Prostate cancer is now the most frequently diagnosed cancer in men. Prostate cancer is latent; many men carry prostate cancer cells without overt signs of disease. It is associated with a high morbidity. Cancer metastasis to bone (late stage) is common and is almost always fatal.

Current treatments include radical prostatectomy, radiation therapy, hormonal ablation and chemotherapy. Unfortunately, in approximately 80% of cases, diagnosis of prostate cancer is established when the disease has already metastasized to the bones, thus limiting the effectiveness of surgical treatments. Hormonal therapy frequently fails with time with the development of hormone-resistant tumor cells. Although chemotherapeutic agents have been used in the treatment of prostate cancer, no single agent has demonstrated superiority over its counterparts, and no drug combination seems particularly effective. The generally drug-resistant, slow-growing nature of most prostate cancers makes them particularly unresponsive to standard chemotherapy.

A major, indeed the overwhelming, obstacle to cancer therapy is the problem of selectivity; that is, the ability to inhibit the multiplication of tumor cells, while leaving unaffected the function of normal cells. For example, in prostate cancer therapy, the therapeutic ratio, or ratio of tumor cell killing to normal cell killing of traditional tumor chemotherapy, is only 1.5:1. Thus, more effective treatment methods and pharmaceutical compositions for therapy and prophylaxis of neoplasia are needed.

Solid tumors frequently contain regions that are poorly vascularized, partly because the tumor cells grow faster than the endothelial cells that make up the blood vessels. Tumor cells can remain viable in such hypoxic conditions and are often refractory to chemotherapy and radiation therapy. In a recent study of cervical cancer, the oxygen status of a tumor was shown to be the single most important prognostic factor, ahead of age of patient, menopausal status, clinical stage, size and histology. Hoeckel et al. (1996) *Semin. Radiat. Oncol.* 6:1-8.

Of particular interest is development of more specific, targeted forms of cancer therapy, especially for cancers that are difficult to treat successfully. In contrast to conventional cancer therapies, which result in relatively non-specific and often serious toxicity, more specific

treatment modalities attempt to inhibit or kill malignant cells selectively while leaving healthy cells intact. Radioresistant and chemoresistant tumors present particular challenges, and there is a need for methods of treating these types of tumors.

One possible treatment approach for many of these cancers is gene therapy, whereby a gene of interest is introduced into the malignant cell. Various viral vectors, including adenoviral vectors, have been developed as vehicles for gene therapy. The virtually exclusive focus in development of adenoviral vectors for gene therapy is use of adenovirus merely as a vehicle for introducing the gene of interest, not as an effector in itself. Replication of adenovirus has been viewed as an undesirable result, largely due to the host immune response. In the treatment of cancer by replication-defective adenoviruses, the host immune response limits the duration of repeat doses at two levels. First, the capsid proteins of the adenovirus delivery vehicle itself are immunogenic. Second, viral late genes are frequently expressed in transduced cells, eliciting cellular immunity. Thus, the ability to repeatedly administer cytokines, tumor suppressor genes, ribozymes, suicide genes, or genes which convert prodrug to an active drug has been limited by the immunogenicity of both the gene transfer vehicle and the viral gene products of the transfer vehicle as well as the transient nature of gene expression.

Use of adenoviral vectors as therapeutic vehicles for cancer has been reported. Some of these approaches utilize tissue (i.e., cell type) specific transcriptional regulatory elements to selectively drive adenoviral replication (and thus cytotoxicity). U.S. Pat. No. 5,698,443; see also WO 95/11984; WO 96/17053; WO 96/34969; WO 98/35028. While useful and promising, there remain other treatment contexts for which tissue specific replication may be insufficient.

Besides cancerous cells, it is often desirable to selectively destroy certain unwanted cells or tissues. Besides surgery, however, which is invasive, there is a dearth of methods available, particularly non-invasive methods, which would allow such selective cytotoxicity and/or suppression.

There is a need for vector constructs that are capable of eliminating essentially all cancerous cells in a minimum number of administrations before specific immunological response against the vector prevents further treatment and which are suitable for use in

specific, focused cancer ablation treatments. There is also a need for an ability to selectively destroy, or impair, unwanted cells, regardless of cell type and/or regardless of anatomical location.

5

SUMMARY OF THE INVENTION

Replication-competent adenoviral vectors specific for cells in a given, or particular, physiological state that permits or induces expression of polynucleotides under transcriptional control of a cell status-specific TRE, and methods for their use are provided. In these replication-competent adenovirus vectors, one or more adenoviral genes is under transcriptional control of an cell status-specific transcriptional regulatory element (TRE). Preferably, the adenoviral gene under transcriptional control of a cell status-specific TRE is one that is essential for adenoviral propagation. A transgene under control of the cell status-specific TRE may also be present. For the adenoviral vectors of the present invention, a cell status-specific TRE is active in a cell existing in a particular, reversible, physiological state, which may be an aberrant physiological state, i.e., a physiological state that deviates from the typical, or normal, physiological state of that same cell when in a non-dividing or regulated dividing state under normal, physiological conditions.

Accordingly, in one aspect, the invention provides an adenovirus vector comprising an adenovirus gene, wherein said adenovirus gene is under transcriptional control of a cell status-specific TRE. In another embodiment, a cell status-specific TRE is human. In another embodiment, a cell status-specific TRE comprises a cell status-specific promoter and enhancer. In yet another embodiment, a cell status-specific TRE is juxtaposed with a cell type-specific TRE, and together the two elements control expression of an adenovirus gene. Thus, the invention provides adenovirus vectors comprising a TRE comprising a cell status-specific TRE and a cell type-specific TRE.

In some embodiments, the adenovirus gene under transcriptional control of a cell status-specific TRE is an adenovirus gene essential for replication. In some embodiments, the

adenoviral gene essential for replication is an early gene. In another embodiment, the early gene is E1A. In another embodiment, the early gene is E1B. In yet another embodiment, both E1A and E1B are under transcriptional control of a cell status-specific TRE. In other embodiments, the adenovirus gene essential for replication is a late gene.

5 In another embodiment, the cell status-specific TRE comprises a hypoxia responsive element. In another embodiment, the cell status-specific TRE comprises the nucleotide sequence of SEQ ID NO:1.

In another embodiment, the cell status-specific TRE comprises a cell cycle-specific TRE. The cell cycle-specific TRE can be derived from the E2F1 5' flanking region. In one
10 embodiment, the cell cycle-specific TRE comprises the nucleotide sequence depicted in SEQ ID NO:2.

In other embodiments, the adenovirus vector can further comprise a transgene, wherein said transgene is under transcriptional control of an cell status-specific TRE. In some
15 embodiments, the transgene is a cytotoxic gene.

In other embodiments, the adenoviral vector comprises an adenoviral gene essential for adenoviral replication under control of a first cell status-specific TRE, and a second adenoviral gene essential for adenoviral replication under control of a second cell status-specific TRE. The first and the second cell status-specific TREs can be identical, substantially identical, or
20 different from, one another.

In other embodiments, the adenoviral vector comprises an adenoviral gene essential for adenoviral replication under control of a first cell status-specific TRE, and a transgene under control of a second cell status-specific TRE. The first and the second cell status-specific TREs can be substantially identical to, or different from, one another.

In other embodiments, the adenovirus vector comprises an adenovirus gene under
25 transcriptional control of a cell status-specific TRE, and a second adenovirus gene under transcriptional control of a cell type-specific TRE. In other embodiments, the adenovirus vector comprises an adenovirus gene under transcriptional control of a cell status-specific TRE, and a transgene under transcriptional control of a cell type-specific TRE.

In another aspect, the invention provides a host cell comprising the adenovirus
30 vector(s) described herein.

In another aspect, the invention provides pharmaceutical compositions comprising an adenovirus vector(s) described herein.

In another aspect, the invention provides kits which contain an adenoviral vector(s) described herein.

5 In another aspect, methods are provided for conferring selective cytotoxicity in target cells (i.e., cells which permit or induce a cell status-specific TRE to function), comprising contacting the cells with an adenovirus vector(s) described herein, whereby the vector enters the cell.

10 Another embodiment of the invention is an adenovirus which replicates preferentially in mammalian cells whose cell status permits or induces the function of a cell status-specific TRE.

In another aspect, methods are provided for propagating an adenovirus specific for mammalian cells whose cell status permits the function of a cell status-specific TRE, said method comprising combining an adenovirus vector(s) described herein with mammalian cells
15 whose cell status permits the function of a cell status-specific TRE, whereby said adenovirus is propagated.

The invention further provides methods of suppressing tumor cell growth, more particularly a target tumor cell (i.e., a tumor cell that permits or induces a cell status-specific TRE to function), comprising contacting a tumor cell with an adenoviral vector of the
20 invention such that the adenoviral vector enters the tumor cell and exhibits selective cytotoxicity for the tumor cell.

In another aspect, methods are provided for detecting cells whose cell status permits the function of a cell status-specific TRE in a biological sample, comprising contacting cells of a biological sample with an adenovirus vector(s) described herein, and detecting replication of
25 the adenovirus vector, if any.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic representation of adenovirus vector CN796, in which the E1A gene is under transcriptional control of an HRE and a PSA-TRE, as described in Example 1.

Figure 2 shows the nucleotide sequence of an HRE from the 5' flanking region of a rat enolase-1 gene (SEQ ID NO:1).

Figure 3 shows the nucleotide sequence of the 5' flanking region of a human E2F1 gene (SEQ ID NO:2). The asterisk indicates the transcription start site.

5 Figure 4 depicts a nucleotide sequence of a prostate-specific antigen TRE.

Figure 5 depicts a nucleotide sequence of a carcinoembryonic antigen TRE.

Figure 6 depicts a nucleotide sequence of a human glandular kallikrein TRE.

Figure 7 depicts a nucleotide sequence of a mucin TRE.

Figure 8 depicts a nucleotide sequence of a rat probasin TRE.

10 Figure 9 depicts a nucleotide sequence and translated amino acid sequence of an adenovirus death protein.

MODES FOR CARRYING OUT THE INVENTION

15 We have discovered and constructed replication-competent adenovirus vectors which contain an adenoviral gene under transcriptional control of a cell status-specific transcriptional response element (TRE) such that the adenovirus gene is transcribed preferentially in cells whose cell status permit the function of the cell status-specific TRE, and have developed methods using these adenovirus vectors. In some preferred embodiments, the adenovirus vectors of this invention comprise at least one adenovirus gene necessary for adenoviral replication, preferably at least one early gene, under the transcriptional control of a TRE specifically regulated by binding of transcriptional factor(s) and/or co-factor(s) necessary for transcription regulated by the cell status-specific TRE. By providing for cell status-specific transcription of at least one adenovirus gene required for replication, the invention provides adenovirus vectors that can be used for specific cytotoxic effects due to selective replication and/or selective transcription. This is especially useful in the cancer context, in which targeted cell killing is desirable. This is also useful for targeted cytotoxic effects in other, non-tumor cells, when selective destruction and/or suppression of these cells is desirable. The vectors can also be useful for detecting the presence of cells whose cell status permits function of a cell status-specific TRE in, for example, an appropriate biological (such as clinical) sample.

25

Further, the adenovirus vector(s) can optionally selectively produce one or more proteins of interest in a target cell by using a cell status-specific TRE.

We have found that adenovirus vectors of the invention replicate and/or express an adenoviral gene operably linked to a cell status-specific TRE preferentially in cells whose status permits the function of a cell status-specific TRE. In contrast to previously- described adenoviral vectors designed to replicate preferentially in specific, differentiated cell types, the adenovirus vectors of the present invention comprise regulatory elements that are not cell type-specific. Rather, they confer cell status-specific adenoviral replication and/or cell status-specific expression of an operably linked adenoviral gene and/or transgene.

The adenovirus vectors of the present invention comprise a cell status-specific TRE which is functional in a cell which exhibits a particular physiological (i.e., environmental or metabolic) characteristic which is reversible and/or progressive. The target cell may exhibit an aberrant physiological state, such as low oxygen tension, or may be subjected to an aberrant environmental condition, such as heat or ionizing radiation, in order for the cell-status TRE to function. The replication preference of these vectors is indicated by comparing the level of replication (i.e., titer) in cells in a requisite physiological state or condition (for example, an aberrant physiological state) to the level of replication in cells not exhibiting the requisite physiological state (for example, under normal physiological conditions). Thus, the invention also uses and takes advantage of what has been considered an undesirable aspect of adenoviral vectors, namely, their replication and possibly concomitant immunogenicity. The probability of runaway infection is significantly reduced due to the cell status-specific requirements for viral replication. Without wishing to be bound by any particular theory, the inventors note that production of adenovirus proteins can serve to activate and/or stimulate the immune system, generally and/or specifically toward target cells producing adenoviral proteins, which can be an important consideration in the cancer context, where patients are often moderately to severely immunocompromised.

The adenovirus vectors of the present invention find particular utility in specific treatment regimens, in which the treatment is highly focused toward, for example, a particular cancer which might otherwise be inoperable or untreatable. An important feature of the

invention is that the vectors are useful in these treatments regardless of the tissue or cell type of the cancer, and yet their cytotoxicity can be targeted to certain locations.

General Techniques

- 5 The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, such as, "Molecular Cloning: A Laboratory Manual", second edition (Sanbrook et al., 1989); "Oligonucleotide Synthesis" (M.J. Gait, ed., 1984); "Animal Cell Culture" (R.I. Freshney, ed., 1987); "Methods in Enzymology" (Academic Press, Inc.); "Handbook of Experimental Immunology" (D.M. Wei & C.C. Blackwell, eds.); "Gene Transfer Vectors for Mammalian Cells" (J.M. Miller & M.P. Calos, eds., 1987); "Current Protocols in Molecular Biology" (F.M. Ausubel et al., eds., 1987); "PCR: The Polymerase Chain Reaction", (Mullis et al., eds., 1994); "Current Protocols in Immunology" (J.E. Coligan et al., eds., 1991).
- 10 For techniques related to adenovirus, see, *inter alia*, Felgner and Ringold (1989) *Nature* 337:387-388; Berkner and Sharp (1983) *Nucl. Acids Res.* 11:6003-6020; Graham (1984) *EMBO J.* 3:2917-2922; Bett et al. (1993) *J. Virology* 67:5911-5921; Bett et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:8802-8806.
- 15

Definitions

- 20 As used herein, a "transcription response element" or "transcriptional regulatory element", or "TRE" is a polynucleotide sequence, preferably a DNA sequence, which increases transcription of an operably linked polynucleotide sequence in a host cell that allows that TRE to function. A TRE can comprise an enhancer and/or a promoter.

- 25 As used herein, the term "cell status-specific TRE" is one that confers transcriptional activation on an operably linked polynucleotide in a cell which allows a cell status-specific TRE to function, i.e., a cell which exhibits a particular physiological condition, including, but not limited to, an aberrant physiological state. "Cell status" thus refers to a given, or particular, physiological state (or condition) of a cell, which is reversible and/or progressive. The physiological state may be generated internally or externally; for example, it may be a

metabolic state (such as low oxygen), or it may be generated due to heat or ionizing radiation. "Cell status" is distinct from a "cell type", which relates to a differentiation state of a cell, which under normal conditions is irreversible. Generally (but not necessarily), as discussed herein, a cell status is embodied in an aberrant physiological state, examples of which are given below.

A "normal cell status" or "normal physiological state" is the status of a cell which exists in normal physiological conditions and which is non-dividing or divides in a regulated manner, i.e., a cell in a normal physiological state.

The terms "aberrant cell status" and "aberrant physiological state", used interchangeably herein, intend a condition of a cell which is a response to, a result of, or is influenced by, an aberrant physiological condition. An aberrant cell status is neither cell type-specific nor tissue type-specific. An aberrant cell status is defined in relation to a cell of the same type which is in a non-dividing/regulated dividing state and under normal physiological conditions.

"Normal physiological conditions" are known to those skilled in the art. These conditions may vary, depending on a cell's location in the body. For example, oxygen tension can vary from tissue to tissue. For *in vitro* analyses for the purposes of determining whether a TRE is responsive to deviations from normal physiological conditions, these conditions generally include an oxygen concentration of about 20% O₂, and a temperature of about 37°C. "Regulated cell division" is a term well understood in the art and refers to the normal mitotic activity of a cell. Those skilled in the art understand that normal mitotic activity varies from cell type to cell type. For example, many terminally differentiated cells in tissues exhibit little or no mitotic activity, while hematopoietic cells are generally mitotically active.

An "aberrant physiological condition" or "aberrant physiological state", as used herein, intends a condition which deviates from normal physiological conditions, and includes, but is not limited to, a physiological condition that is characterized by alterations in oxygen concentration, such as hypoxic conditions; temperatures which deviate from physiological temperatures; a condition that triggers apoptosis; radiation, including ionizing radiation and UV irradiation; de-regulated cell division, resulting for example, from a lack of, or insufficient amounts of, or inactivity of, a factor which controls cell division, such as, for example,

retinoblastoma protein (Rb); variations in timing of cell cycle; infection with a pathogen; exposure to a chemical substance; or a combination of the above-listed conditions. Another example is a mutation that could, or does, exist in any cell type, i.e., its existence does not depend on, or is not related to, the differentiation state of the cell.

5 A "target cell", as used herein, is one that permits or induces the function of a cell status-specific TRE such that it effects transcriptional activation of an operably linked polynucleotide. A target cell is one which exhibits a requisite physiological (or environmental) state, which may be an aberrant physiological state. Preferably, a target cell is a mammalian cell, preferably a human cell. A target cell may or may not be neoplastic. By
10 transcriptional activation, it is intended that transcription is increased in the target cell above the levels in a control cell (e.g., a that cell when not exhibiting a requisite physiological state (generally a normal physiological state) by at least about 2 fold, preferably at least about 5 fold, preferably at least about 10 fold, more preferably at least about 20 fold, more preferably at least about 50 fold, more preferably at least about 100 fold, more preferably at least about
15 200 fold, even more preferably at least about 400 fold to about 500 fold, even more preferably at least about 1000 fold. The normal levels are generally the level of activity (if any) in a cell as tested under conditions that activate the cell status-specific TRE, or the level of activity (if any) of a reporter construct lacking a cell status-specific TRE as measured in a cell exhibiting the requisite physiological condition.

20 A "functionally-preserved" variant of a cell status-specific TRE is a cell status-specific TRE which differs from another cell status-specific TRE, but still retains cell status cell-specific transcription activity. The difference in an cell status-specific TRE can be due to differences in linear sequence, arising from, for example, single base mutation(s), addition(s), deletion(s), and/or modification(s) of the bases. The difference can also arise from changes in
25 the sugar(s), and/or linkage(s) between the bases of a cell status-specific TRE.

An "adenovirus vector" or "adenoviral vector" (used interchangeably) comprises a polynucleotide construct of the invention. A polynucleotide construct of this invention may be in any of several forms, including, but not limited to, DNA, DNA encapsulated in an adenovirus coat, DNA packaged in another viral or viral-like form (such as herpes simplex, and AAV), DNA encapsulated in liposomes, DNA complexed with polylysine, complexed
30

with synthetic polycationic molecules, conjugated with transferrin, and complexed with compounds such as PEG to immunologically "mask" the molecule and/or increase half-life, and conjugated to a nonviral protein. Preferably, the polynucleotide is DNA. As used herein, "DNA" includes not only bases A, T, C, and G, but also includes any of their analogs or modified forms of these bases, such as methylated nucleotides, internucleotide modifications such as uncharged linkages and thioates, use of sugar analogs, and modified and/or alternative backbone structures, such as polyamides. For purposes of this invention, adenovirus vectors are replication-competent in a target cell.

The terms "polynucleotide" and "nucleic acid", used interchangeably herein, refer to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. These terms include a single-, double- or triple-stranded DNA, genomic DNA, cDNA, RNA, DNA-RNA hybrid, or a polymer comprising purine and pyrimidine bases, or other natural, chemically, biochemically modified, non-natural or derivatized nucleotide bases. The backbone of the polynucleotide can comprise sugars and phosphate groups (as may typically be found in RNA or DNA), or modified or substituted sugar or phosphate groups. Alternatively, the backbone of the polynucleotide can comprise a polymer of synthetic subunits such as phosphoramidates and thus can be a oligodeoxynucleoside phosphoramidate (P-NH₂) or a mixed phosphoramidate- phosphodiester oligomer. Peyrottes et al. (1996) *Nucleic Acids Res.* 24: 1841-8; Chaturvedi et al. (1996) *Nucleic Acids Res.* 24: 2318-23; Schultz et al. (1996) *Nucleic Acids Res.* 24: 2966-73. A phosphorothiate linkage can be used in place of a phosphodiester linkage. Braun et al. (1988) *J. Immunol.* 141: 2084-9; Latimer et al. (1995) *Mol. Immunol.* 32: 1057-1064. In addition, a double-stranded polynucleotide can be obtained from the single stranded polynucleotide product of chemical synthesis either by synthesizing the complementary strand and annealing the strands under appropriate conditions, or by synthesizing the complementary strand de novo using a DNA polymerase with an appropriate primer.

The following are non-limiting examples of polynucleotides: a gene or gene fragment, exons, introns, mRNA, tRNA, rRNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers. A polynucleotide may comprise modified

nucleotides, such as methylated nucleotides and nucleotide analogs, uracyl, other sugars and linking groups such as fluororibose and thioate, and nucleotide branches. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component.

5 Other types of modifications included in this definition are caps, substitution of one or more of the naturally occurring nucleotides with an analog, and introduction of means for attaching the polynucleotide to proteins, metal ions, labeling components, other polynucleotides, or a solid support. Preferably, the polynucleotide is DNA. As used herein, "DNA" includes not only bases A, T, C, and G, but also includes any of their analogs or modified forms of these bases,
10 such as methylated nucleotides, internucleotide modifications such as uncharged linkages and thioates, use of sugar analogs, and modified and/or alternative backbone structures, such as polyamides.

A polynucleotide or polynucleotide region has a certain percentage (for example, 80%, 85%, 90%, or 95%) of "sequence identity" to another sequence means that, when aligned, that
15 percentage of bases are the same in comparing the two sequences. This alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example those described in *Current Protocols in Molecular Biology* (F.M. Ausubel et al., eds., 1987) Supplement 30, section 7.7.18, Table 7.7.1. A preferred alignment program is ALIGN Plus (Scientific and Educational Software, Pennsylvania), preferably using default
20 parameters.

"Under transcriptional control" is a term well-understood in the art and indicates that transcription of a polynucleotide sequence, usually a DNA sequence, depends on its being operably (operatively) linked to an element which contributes to the initiation of, or promotes, transcription. "Operably linked" refers to a juxtaposition wherein the elements are in an
25 arrangement allowing them to function.

"Replication" and "propagation" are used interchangeably and refer to the ability of a polynucleotide construct of the invention to reproduce, or proliferate. This term is well understood in the art. For purposes of this invention, replication involves production of adenovirus proteins and is generally directed to reproduction of adenovirus. Replication can

be measured using assays standard in the art and described herein, such as a burst assay, plaque assay, or a one-step growth curve assay.

As used herein, "cytotoxicity" is a term well understood in the art and refers to a state in which a cell's usual biochemical or biological activities are compromised (i.e., inhibited). These activities include, but are not limited to, metabolism; cellular replication; DNA
5 replication; transcription; translation; uptake of molecules. "Cytotoxicity" includes cell death and/or cytolysis. Assays are known in the art which indicate cytotoxicity, such as dye exclusion, ³H-thymidine uptake, and plaque assays.

The term "selective cytotoxicity", as used herein, refers to the cytotoxicity conferred by an adenovirus vector of the present invention on a cell which allows or induces a cell status-specific TRE to function (a target cell) when compared to the cytotoxicity conferred by an
10 adenoviral vector of the present invention on a cell which does not allow a cell status-specific TRE to function (a non-target cell). Such cytotoxicity may be measured, for example, by plaque assays, by reduction or stabilization in size of a tumor comprising target cells, or the
15 reduction or stabilization of serum levels of a marker characteristic of the tumor cells, or a tissue-specific marker, e.g., a cancer marker, such as prostate specific antigen.

In the context of adenovirus, a "heterologous polynucleotide" or "heterologous gene" or "transgene" is any polynucleotide or gene that is not present in wild-type adenovirus. Preferably, the transgene will also not be expressed or present in the target cell prior to
20 introduction by the adenovirus vector. Examples of preferred transgenes are provided below.

In the context of adenovirus, a "heterologous" promoter or enhancer is one which is not associated with or derived from an adenovirus gene.

In the context of adenovirus, an "endogenous" promoter, enhancer, or TRE is native to or derived from adenovirus.

In the context of a cell status-specific TRE, a "heterologous" promoter or enhancer is one which is not normally associated in a cell with or derived from a cell status-specific TRE. Examples of a heterologous promoter or enhancer are the albumin promoter or enhancer and other viral promoters and enhancers, such as SV40, or cell type-specific TREs such as a
25 prostate-specific TRE.

A "cell type-specific TRE" is preferentially functional in a specific type of cell relative to other types of cells. In contrast to cell status, "cell type" is a reflection of a differentiation state of a cell which is irreversible. For example, a prostate-specific antigen is expressed in prostate cells, but is not substantially expressed in other cell types such as hepatocytes, astrocytes, cardiocytes, lymphocytes, etc. Generally, a cell type-specific TRE is active in only one cell type. When a cell type-specific TRE is active in more than one cell type, its activity is restricted to a limited number of cell types, i.e., it is not active in all cell types. A cell type-specific TRE may or may not be tumor cell specific.

"Suppressing" tumor growth indicates a growth state that is curtailed when compared to growth without contact with, i.e., transfection by, an adenoviral vector described herein. Tumor cell growth can be assessed by any means known in the art, including, but not limited to, measuring tumor size, determining whether tumor cells are proliferating using a ³H-thymidine incorporation assay, or counting tumor cells. "Suppressing" tumor cell growth means any or all of the following states: slowing, delaying, and stopping tumor growth, as well as tumor shrinkage.

As used herein, the terms "neoplastic cells", "neoplasia", "tumor", "tumor cells", "cancer" and "cancer cells", (used interchangeably) refer to cells which exhibit relatively autonomous growth, so that they exhibit an aberrant growth phenotype characterized by a significant loss of control of cell proliferation (i.e., de-regulated cell division). Neoplastic cells can be malignant or benign.

A "host cell" includes an individual cell or cell culture which can be or has been a recipient of an adenoviral vector(s) of this invention. Host cells include progeny of a single host cell, and the progeny may not necessarily be completely identical (in morphology or in total DNA complement) to the original parent cell due to natural, accidental, or deliberate mutation and/or change. A host cell includes cells transfected or infected *in vivo* or *in vitro* with an adenoviral vector of this invention.

"Replication" and "propagation" are used interchangeably and refer to the ability of an adenovirus vector of the invention to reproduce or proliferate. These terms are well understood in the art. For purposes of this invention, replication involves production of adenovirus proteins and is generally directed to reproduction of adenovirus. Replication can

be measured using assays standard in the art and described herein, such as a burst assay or plaque assay. "Replication" and "propagation" include any activity directly or indirectly involved in the process of virus manufacture, including, but not limited to, viral gene expression; production of viral proteins, nucleic acids or other components; packaging of viral components into complete viruses; and cell lysis.

An "ADP coding sequence" is a polynucleotide that encodes ADP or a functional fragment thereof. In the context of ADP, a "functional fragment" of ADP is one that exhibits cytotoxic activity, especially cell lysis, with respect to adenoviral replication. Ways to measure cytotoxic activity are known in the art and are described herein.

A polynucleotide that "encodes" an ADP polypeptide is one that can be transcribed and/or translated to produce an ADP polypeptide or a fragment thereof. The anti-sense strand of such a polynucleotide is also said to encode the sequence.

An "ADP polypeptide" is a polypeptide containing at least a portion, or region, of the amino acid sequence of an ADP (see, for example, SEQ ID NO:5), and which displays a function associated with ADP, particularly cytotoxicity, more particularly, cell lysis. As discussed herein, these functions can be measured using techniques known in the art. It is understood that certain sequence variations may be used, due to, for example, conservative amino acid substitutions, which may provide ADP polypeptides.

A polynucleotide sequence that is "depicted in" a SEQ ID NO means that the sequence is present as an identical contiguous sequence in the SEQ ID NO. The term encompasses portions, or regions of the SEQ ID NO as well as the entire sequence contained within the SEQ ID NO.

A "biological sample" encompasses a variety of sample types obtained from an individual and can be used in a diagnostic or monitoring assay. The definition encompasses blood and other liquid samples of biological origin, solid tissue samples such as a biopsy specimen or tissue cultures or cells derived therefrom, and the progeny thereof. The definition also includes samples that have been manipulated in any way after their procurement, such as by treatment with reagents, solubilization, or enrichment for certain components, such as proteins or polynucleotides. The term "biological sample" encompasses a clinical sample, and

also includes cells in culture, cell supernatants, cell lysates, serum, plasma, biological fluid, and tissue samples.

An "individual" is a vertebrate, preferably a mammal, more preferably a human. Mammals include, but are not limited to, farm animals, sport animals, rodents, primates, and
5 pets.

An "effective amount" is an amount sufficient to effect beneficial or desired results, which may include clinical results. An effective amount can be administered in one or more administrations. For purposes of this invention, an effective amount of an adenoviral vector is an amount that is sufficient to palliate, ameliorate, stabilize, reverse, slow or delay the
10 progression of the disease state.

Adenoviral vectors comprising a cell status-specific TRE

The present invention provides adenoviral vector constructs which comprise an adenovirus gene under transcriptional control of a cell status-specific TRE. Preferably, the adenovirus gene contributes to cytotoxicity (whether direct and/or indirect), more preferably one that contributes to or causes cell death, even more preferably is essential for adenoviral
15 replication. Examples of a gene that contributes to cytotoxicity include, but are not limited to, adenovirus death protein (ADP; discussed below). When the adenovirus vector(s) is selectively (i.e., preferentially) replication competent for propagation in target cells, i.e., cells which permit or induce a cell-status TRE to function, these cells will be preferentially killed
20 upon adenoviral proliferation. Once the target cells are destroyed due to selective cytotoxic and/or cytolytic replication, the adenovirus vector replication is significantly reduced, thus lessening the probability of runaway infection and undesirable bystander effects. *In vitro* cultures may be retained to monitor the mixture (such as, for example, a biopsy or other appropriate biological sample) for occurrence (i.e., presence) and/or recurrence of the target
25 cell, e.g., a neoplastic cell or other undesired cell. To further ensure cytotoxicity, one or more transgenes having a cytotoxic effect may also be present and under selective transcriptional control. In this embodiment, one may provide higher confidence that the target cells will be destroyed. Additionally, or alternatively, an adenovirus gene that contributes to cytotoxicity

and/or cell death (such as ADP) may be included in the adenoviral vector, either free of, or under, selective transcriptional control.

Cell status-specific TREs for use in the adenoviral vectors of the present invention can be derived from any species, preferably a mammal. A number of genes have been described which are expressed in response to, or in association with, a cell status. Any of these cell status-associated genes may be used to generate a cell status-specific TRE.

An example of a cell status is cell cycle. An exemplary gene whose expression is associated with cell cycle is E2F-1, a ubiquitously expressed, growth-regulated gene, which exhibits peak transcriptional activity in S phase. Johnson et al. (1994) *Genes Dev.* 8:1514-1525. The RB protein, as well as other members of the RB family, form specific complexes with E2F-1, thereby inhibiting its ability to activate transcription. Thus, E2F-1-responsive promoters are down-regulated by RB. Many tumor cells have disrupted RB function, which can lead to de-repression of E2F-1-responsive promoters, and, in turn, de-regulated cell division.

Accordingly, in one embodiment, the invention provides an adenoviral vector in which an adenoviral gene (preferably a gene necessary for replication) is under transcriptional control of a cell status-specific TRE, wherein the cell status-specific TRE comprises a cell cycle-activated, or cell-cycle specific, TRE. In one embodiment, the cell cycle-activated TRE is an E2F1 TRE. In one embodiment, this TRE comprises the sequence depicted in Figure 3 and SEQ ID NO:2.

Another group of genes which are regulated by cell status are those whose expression is increased in response to hypoxic conditions. Bunn and Poyton (1996) *Physiol. Rev.* 76:839-885; Dachs and Stratford (1996) *Br. J. Cancer* 74:5126-5132; Guillemin and Krasnow (1997) *Cell* 89:9-12. Many tumors have insufficient blood supply, due in part to the fact that tumor cells typically grow faster than the endothelial cells that make up the blood vessels, resulting in areas of hypoxia in the tumor. Folkman (1989) *J. Natl. Cancer Inst.* 82:4-6; and Kallinowski (1996) *The Cancer J.* 9:37-40. An important mediator of hypoxic responses is the transcriptional complex HIF-1, or hypoxia inducible factor-1, which interacts with a hypoxia-responsive element (HRE) in the regulatory regions of several genes, including vascular endothelial growth factor, and several genes encoding glycolytic enzymes, including enolase-

1. Murine HRE sequences have been identified and characterized. Firth et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:6496-6500. An HRE from a rat enolase-1 promoter is described in Jiang et al. (1997) *Cancer Res.* 57:5328-5335. An HRE from a rat enolase-1 promoter is depicted in Figure 2 and given as SEQ ID NO:1.

5 Accordingly, in one embodiment, an adenovirus vector comprises an adenovirus gene, preferably an adenoviral gene essential for replication, under transcriptional control of a cell status-specific TRE comprising an HRE. In one embodiment, the cell status-specific TRE comprises the HRE depicted in Figure 2 and SEQ ID NO:1.

10 Other cell status-specific TREs include heat-inducible (i.e., heat shock) promoters, and promoters responsive to radiation exposure, including ionizing radiation and UV radiation. For example, the promoter region of the early growth response-1 (Egr-1) gene contains an element(s) inducible by ionizing radiation. Hallahan et al. (1995) *Nat. Med.* 1:786-791; and Tsai-Morris et al. (1988) *Nucl. Acids. Res.* 16:8835-8846. Heat-inducible promoters, including heat-inducible elements, have been described. See, for example Welsh (1990) in "Stress
15 Proteins in Biology and Medicine", Morimoto, Tisseries, and Georgopoulos, eds. Cold Spring Harbor Laboratory Press; and Perisic et al. (1989) *Cell* 59:797-806. Accordingly, in some embodiments, the cell status-specific TRE comprises an element(s) responsive to ionizing radiation. In one embodiment, this TRE comprises a 5' flanking sequence of an Egr-1 gene. In other embodiments, the cell status-specific TRE comprises a heat shock responsive, or heat-
20 inducible, element.

 A cell status-specific TRE can also comprise multimers. For example, an HRE can comprise a tandem series of at least two, at least three, at least four, or at least five hypoxia-responsive elements. These multimers may also contain heterologous promoter and/or enhancer sequences.

25 A cell status-specific TRE may or may not lack a silencer. The presence of a silencer (i.e., a negative regulatory element) may assist in shutting off transcription (and thus replication) in non-permissive cells (i.e., cell in a normal cell state). Thus, presence of a silencer may confer enhanced cell status-specific replication by more effectively preventing adenoviral vector replication in non-target cells. Alternatively, lack of a silencer may assist in

effecting replication in target cells, thus conferring enhanced cell status-specific replication due to more effective replication in target cells.

In other embodiments, the adenoviral vector comprises an adenoviral gene essential for adenoviral replication under control of a first cell status-specific TRE, and a second adenoviral gene essential for adenoviral replication under control of a second cell status-specific TRE. The first and the second cell status-specific TREs may or may not be identical, and may or may not be substantially identical to one another. By "substantially identical" is meant a requisite degree of sequence identity between the two TREs. The degree of sequence identity between these TREs is at least about 80%, preferably at least about 85%, more preferably at least about 90%, even more preferably at least about 95%, even more preferably at least about 98%, and most preferably 100%. Sequence identity can be determined by a sequence comparison using, i.e., sequence alignment programs that are known in the art, such as those described in *Current Protocols in Molecular Biology* (F.M. Ausubel et al., eds., 1987) Supplement 30, section 7.7.18, Table 7.7.1. A preferred alignment program is ALIGN Plus (Scientific and Educational Software, Pennsylvania), preferably using default parameters. Alternatively, hybridization under stringent conditions can also indicate degree of sequence identity. Stringent conditions are known in the art; an example of a stringent condition is 80°C (or higher temperature) and 6 X SSC (or less concentrated SSC). Other hybridization conditions and parameters (in order of increasing stringency) are: incubation temperatures of 25°C, 37°C, 50°C, and 68°C; buffer concentrations of 10 X SSC, 6 X SSC, 1 X SSC, 0.1 X SSC (where 1 X SSC is 0.15 M NaCl and 15 mM citrate buffer) and their equivalents using other buffer systems; formamide concentrations of 0%, 25%, 50%, and 75%; incubation times from about 24 hours about 5 minutes; 1, 2, or more washing steps; wash incubation times of 1, 2, or 15 minutes; and wash solutions of 6 X SSC, 1 X SSC, 0.1 X SSC, or deionized water.

Adenoviral constructs in which the first and second cell status-specific TREs are identical or substantially identical, particularly if these TREs control transcription of early genes (such as E1A and E1B), may display an instability which may be desirable in certain contexts, such as when an automatic "self-destruction" property can shut down the virus, thereby controlling the degree of propagation. Accordingly, in some embodiments, the first and second cell status-specific TRE, or the first and second TRE (one of which is a cell-status-

specific TRE) are sufficiently identical to confer instability when compared to two TREs which are less identical with respect to each other (i.e., have more sequence divergence or dissimilarity). Preferred embodiments are those in which the two TREs control E1A and E1B respectively. "Instability" means that the structural integrity of the adenoviral vectors is not preserved as the virus replicates in cells, and can be measured using standard methods in the art, such as Southern analysis. In other embodiments, the first and second TREs are sufficiently divergent and/or placed in the vector such that the vector is stable (i.e., the structural integrity of the adenoviral vector is preserved).

In other embodiments, the adenoviral vector comprises an adenoviral gene essential for adenoviral replication under control of a first cell status-specific TRE, and a transgene under control of a second cell status-specific TRE. The first and the second cell status-specific TREs may or may not be substantially identical to one another.

In some embodiments, a cell status-specific TRE can be juxtaposed with another TRE, such as a different cell status-specific TRE, or, alternatively, a cell type-specific TRE. "Juxtaposed" means a cell status-specific TRE and the second TRE transcriptionally control the same gene, or at least are proximate with respect to the same gene. For these embodiments, the cell status-specific TRE and the second TRE may be in any of a number of configurations, including, but not limited to, (a) next to each other (i.e., abutting); (b) both 5' to the gene that is transcriptionally controlled (i.e., may have intervening sequences between them); (c) one TRE 5' and the other TRE 3' to the gene. For example, as described in Example 1 and shown in Figure 1, a cell type-specific TRE can be juxtaposed with a cell status-specific TRE to control transcription of an operably linked adenoviral gene. Such "composite" TREs can be used to confer both cell status- and cell type-specific expression of an operably linked polynucleotide, and thus may confer significantly greater specificity and/or efficacy. Examples of cell type-specific TREs are provided below. Alternatively, "composite" TREs can be used to confer different, and possibly synergistic, cell status specificity. For example, a composite TRE could confer specificity to hypoxia and heat shock.

Example 1 provides a description of an adenovirus construct in which a composite TRE upstream of E1A consisting of an HRE and a prostate-specific TRE, PSA-TRE (which consists of enhancer sequences -5322 to -3738 fused to PSA promoter sequence -541 to +12; see U.S.

Pat. Nos. 5,871,726; 5,648,478). Accordingly, in some embodiments, the invention provides an adenovirus vector comprising an adenovirus gene essential for replication, preferably an early gene, preferably E1A or E1B, under transcriptional control of a TRE comprising an HRE (preferably comprising or consisting of the 67-base fragment depicted in SEQ ID NO:1) and a prostate cell specific TRE, preferably comprising a PSA enhancer (preferably -5322 to -3738; or about 503 to about 2086 of SEQ ID NO:3 (bases about 503 to about 2086 of Figure 4), and a promoter, preferably comprising a PSA enhancer and a PSA promoter (about 5285 to about 5836 of SEQ ID NO:3).

As is readily appreciated by one skilled in the art, a cell status-specific TRE is a polynucleotide sequence, and, as such, can exhibit function over a variety of sequence permutations. Methods of nucleotide substitution, addition, and deletion are known in the art, and readily available functional assays (such as the CAT or luciferase reporter gene assay) allow one of ordinary skill to determine whether a sequence variant exhibits requisite cell status-specific transcription function. Hence, the invention also includes functionally-preserved variants of the nucleic acid sequences disclosed herein, which include nucleic acid substitutions, additions, and/or deletions. While not wishing to be bound by a single theory, the inventors note that it is possible that certain modifications will result in modulated resultant expression levels, including enhanced expression levels. Achievement of modulated resultant expression levels, preferably enhanced expression levels, may be especially desirable in the case of certain, more aggressive forms of cancer, or when a more rapid and/or aggressive pattern of cell killing is warranted (due to an immunocompromised condition of the individual, for example).

As an example of how cell status-specific TRE activity can be determined, a polynucleotide sequence or set of such sequences can be generated using methods known in the art, such as chemical synthesis, site-directed mutagenesis, PCR, and/or recombinant methods. The sequence(s) to be tested is inserted into a vector containing an appropriate reporter gene, including, but not limited to, chloramphenicol acetyl transferase (CAT), β -galactosidase (encoded by the lacZ gene), luciferase (encoded by the luc gene), green fluorescent protein, alkaline phosphatase, and horse radish peroxidase. Such vectors and assays are readily available, from, inter alia, commercial sources. Plasmids thus constructed

are transfected into a suitable host cell to test for expression of the reporter gene as controlled by the putative cell status-specific TRE using transfection methods known in the art, such as calcium phosphate precipitation, electroporation, liposomes (lipofection), and DEAE-dextran. Suitable host cells include any cell type, including but not limited to, Hep3B, Hep G2, HuH7, HuH1/C12, LNCaP, HBL-100, Chang liver cells, MCF-7, HLF, HLE, 3T3, HUVEC, and HeLa. Host cells transfected with the TRE-reporter gene construct to be tested are subjected to conditions which result in a change in cell status (for example, one which result in an aberrant physiological state). The same cells not subjected to these conditions, i.e., which are under normal physiological conditions and therefore in a normal physiological state, serve as controls. Results are obtained by measuring the level of expression of the reporter gene using standard assays. Comparison of expression between cells in a particular state and control indicates presence or absence of transcriptional activation. "Transcriptional activation" has been defined above.

Comparisons between or among various cell status-specific TREs can be assessed, for example, by measuring and comparing levels of expression within a single cell line under normal and aberrant physiological conditions. These comparisons may also be made by measuring and comparing levels of expression within a single cell line subjected to reversible environmental conditions (such as heat) and cells not subjected to such conditions. It is understood that absolute transcriptional activity of an cell status-specific TRE will depend on several factors, such as the nature of the target cell, delivery mode and form of the cell status-specific TRE, and the coding sequence that is to be selectively transcriptionally activated. To compensate for various plasmid sizes used, activities can be expressed as relative activity per mole of transfected plasmid. Alternatively, the level of transcription (i.e., mRNA) can be measured using standard Northern analysis and hybridization techniques. Levels of transfection (i.e., transfection efficiencies) are measured by co-transfecting a plasmid encoding a different reporter gene under control of a different TRE, such as the cytomegalovirus (CMV) immediate early promoter. This analysis can also indicate negative regulatory regions, i.e., silencers.

As an example of how hypoxia induction can be measured, one can use an assay such as that described in Jiang et al. (1997) *Cancer Research* 57:5328-5335 or Dachs et al. (1997)

Nature Med. 3:515-520. For example, a construct comprising a putative HRE, or multiple tandem copies thereof, together with a minimal promoter element, operably linked and controlling transcription of a polynucleotide which encodes a protein which is detectable or can be used to give a detectable signal, is introduced into host cells. The host cells are then subjected to conditions of normoxia (e.g., 20% O₂), and varying degrees of hypoxia, such as 5%, 2%, 1%, 0.1%, or less, O₂. The expression product of the operably linked polynucleotide (reporter gene) is then measured.

Alternatively a putative cell status-specific TRE can be assessed for its ability to confer adenoviral replication preference for cells exhibiting the requisite physiological state, such as heat or ionizing radiation. For this assay, constructs containing an adenovirus gene essential to replication operably linked to a putative cell status-specific TRE are transfected into cells which exhibit the requisite physiological state. Viral replication in those cells is compared, for example, to viral replication by the construct in cells under normal physiological conditions (i.e., not exhibiting the requisite physiological state).

Any of the various serotypes of adenovirus can be used, such as Ad2, Ad5, Ad12 and Ad40. For purposes of illustration, serotype Ad5 will be exemplified herein.

When a cell status-specific TRE is used with an adenovirus gene that is essential for propagation replication competence is preferentially achievable in the target cell expressing cell status. Preferably, the gene is an early gene, such as E1A, E1B, E2, or E4. (E3 is not essential for viral replication.) More preferably, the early gene under cell status-TRE control is E1A and/or E1B. More than one early gene can be placed under control of an cell status-specific TRE. Example 1 provides a more detailed description of such constructs.

The E1A gene is expressed immediately after viral infection (0-2 hours) and before any other viral genes. E1A protein acts as a *trans*-acting positive-acting transcriptional regulatory factor, and is required for the expression of the other early viral genes E1B, E2, E3, E4, and the promoter-proximal major late genes. Despite the nomenclature, the promoter proximal genes driven by the major late promoter are expressed during early times after Ad5 infection. Flint (1982) *Biochem. Biophys. Acta* 651:175-208; Flint (1986) *Advances Virus Research* 31:169-228; Grand (1987) *Biochem. J.* 241:25-38. In the absence of a functional E1A gene, viral infection does not proceed, because the gene products necessary for viral DNA

replication are not produced. Nevins (1989) *Adv. Virus Res.* 31:35-81. The transcription start site of Ad5 E1A is at 498 and the ATG start site of the E1A protein is at 560 in the virus genome.

The E1B protein functions *in trans* and is necessary for transport of late mRNA from the nucleus to the cytoplasm. Defects in E1B expression result in poor expression of late viral proteins and an inability to shut off host cell protein synthesis. The promoter of E1B has been implicated as the defining element of difference in the host range of Ad40 and Ad5: clinically Ad40 is an enterovirus, whereas Ad5 causes acute conjunctivitis. Bailey, Mackay et al. (1993) *Virology* 193:631; Bailey et al. (1994) *Virology* 202:695-706). The E1B promoter of Ad5 consists of a single high-affinity recognition site for Spl and a TATA box.

The E2 region of adenovirus codes for proteins related to replication of the adenoviral genome, including the 72 kDa DNA-binding protein, the 80 kD precursor terminal protein and the viral DNA polymerase. The E2 region of Ad5 is transcribed in a rightward orientation from two promoters, termed E2 early and E2 late, mapping at 76.0 and 72.0 map units, respectively. While the E2 late promoter is transiently active during late stages of infection and is independent of the E1A transactivator protein, the E2 early promoter is crucial during the early phases of viral replication.

The E2 late promoter overlaps with the coding sequences of a gene encoded by the counterstrand and is therefore not amenable to genetic manipulation. However, the E2 early promoter overlaps only for a few base pairs with sequences coding for a 33 kD protein on the counterstrand. Notably, the SpeI restriction site (Ad5 position 27082) is part of the stop codon for the above mentioned 33 kD protein and conveniently separates the major E2 early transcription initiation site and TATA-binding protein site from the upstream transcription factor binding sites E2F and ATF. Therefore, insertion of a cell status-TRE having SpeI ends into the SpeI site in the +-strand would disrupt the endogenous E2 early promoter of Ad5 and should allow cell status-restricted expression of E2 transcripts.

The E4 gene has a number of transcription products. The E4 region codes for two polypeptides which are responsible for stimulating the replication of viral genomic DNA and for stimulating late gene expression. The protein products of open reading frames (ORFS) 3 and 6 can both perform these functions by binding the 55kD protein from E1B and

heterodimers of E2F-1 and DP-1. The ORF 6 protein requires interaction with the E1B 55kD protein for activity while the ORF 3 protein does not. In the absence of functional protein from ORF 3 and ORF 6, plaques are produced with an efficiency less than 10^{-6} that of wild type virus. To further restrict viral replication to cells exhibiting a requisite physiological condition or state, E4 ORFs 1-3 can be deleted, making viral DNA replication and late gene synthesis dependent on E4 ORF 6 protein. By combining such a mutant with sequences in which the E1B region is regulated by a cell status-specific TRE, a virus can be obtained in which both the E1B function and E4 function are dependent on a cell status-specific TRE driving E1B.

The major late genes relevant to the subject invention are genes L1, L2, L3, L4, and L5 which encode proteins of the adenovirus virion. All of these genes (typically coding for structural proteins) are probably required for adenoviral replication. The late genes are all under the control of the major late promoter (MLP), which is located in Ad5 at +5986 to +6048.

In addition to conferring selective cytotoxic and/or cytolytic activity by virtue of preferential replication competence in cells exhibiting a requisite physiological state (for example, an aberrant physiological state such as low oxygen conditions), the adenovirus vectors of this invention can further include a heterologous gene (transgene) under the control of a cell status-specific TRE. In this way, various genetic capabilities may be introduced into target cells, particularly cancer cells. For example, in certain instances, it may be desirable to enhance the degree and/or rate of cytotoxic activity, due to, for example, the relatively refractory nature or particular aggressiveness of the cancerous target cell. This could be accomplished by coupling the cell status-specific replicative cytotoxic activity with cell-specific expression of, for example, HSV-tk and/or cytosine deaminase (cd), which renders cells capable of metabolizing 5-fluorocytosine (5-FC) to the chemotherapeutic agent 5-fluorouracil (5-FU). Using these types of transgenes may also confer a bystander effect.

Other desirable transgenes that may be introduced via an adenovirus vector(s) include genes encoding cytotoxic proteins, such as the A chains of diphtheria toxin, ricin or abrin (Palmiter et al. (1987) *Cell* 50: 435; Maxwell et al. (1987) *Mol. Cell. Biol.* 7: 1576; Behringer et al. (1988) *Genes Dev.* 2: 453; Messing et al. (1992) *Neuron* 8: 507; Piatak et al. (1988) *J.*

Biol. Chem. 263: 4937; Lamb et al. (1985) *Eur. J. Biochem.* 148: 265; Frankel et al. (1989) *Mol. Cell. Biol.* 9: 415), genes encoding a factor capable of initiating apoptosis, sequences encoding antisense transcripts or ribozymes, which among other capabilities may be directed to mRNAs encoding proteins essential for proliferation, such as structural proteins, or transcription factors; viral or other pathogenic proteins, where the pathogen proliferates intracellularly; genes that encode an engineered cytoplasmic variant of a nuclease (e.g. RNase A) or protease (e.g. aprotinin, papain, proteinase K, carboxypeptidase, etc.), or encode the Fas gene, and the like. Other genes of interest include cytokines, antigens, transmembrane proteins, and the like, such as IL-1, -2, -6, -12, GM-CSF, G-CSF, M-CSF, IFN- α , - β , - γ , TNF- α , - β , TGF- α , - β , NGF, and the like. The positive effector genes could be used in an earlier phase, followed by cytotoxic activity due to replication.

In one embodiment, the adenovirus death protein (ADP), encoded within the E3 region, is maintained in the adenovirus vector. The ADP gene, under control of the major late promoter (MLP), appears to code for a protein (ADP) that is important in expediting host cell lysis. Tollefson et al. (1996) *J. Virol.* 70(4):2296; Tollefson et al. (1992) *J. Virol.* 66(6):3633. Thus, adenoviral vectors containing the ADP gene may render the adenoviral vector more potent, making possible more effective treatment and/or a lower dosage requirement.

Accordingly, the invention provides an adenoviral vector as described herein that further includes a polynucleotide sequence encoding an ADP. A DNA sequence encoding an ADP and the amino acid sequence of an ADP are depicted Figure 9. Briefly, an ADP coding sequence is obtained preferably from Ad2 (since this is the strain in which ADP has been more fully characterized) using techniques known in the art, such as PCR. Preferably, the Y leader (which is an important sequence for correct expression of late genes) is also obtained and ligated to the ADP coding sequence. The ADP coding sequence (with or without the Y leader) can then be introduced into the adenoviral genome, for example, in the E3 region (where the ADP coding sequence will be driven by the MLP). The ADP coding sequence could also be inserted in other locations of the adenovirus genome, such as the E4 region. Alternatively, the ADP coding sequence could be operably linked to a heterologous promoter (with or without enhancer(s)), including, but not limited to, another viral promoter, a cell status-specific TRE

such as a hypoxia responsive element, or a cell type-specific TRE such as those derived from carcinoembryonic antigen (CEA), mucin, and rat probasin genes.

Adenoviral vectors of the invention further comprising a cell type specific element

In addition to conferring selective cytotoxic and/or cytolytic activity by virtue of preferential replication competence and/or by preferential transcription of a gene encoding a cytotoxic factor in cells exhibiting a requisite physiological state, the adenovirus vectors of this invention can further include an adenovirus gene and/or a heterologous gene (transgene) under the control of a cell type-specific TRE. In this way, cytotoxicity is further limited to a particular cell type.

For example, TREs that function preferentially in prostate cells include, but are not limited to, TREs derived from the prostate-specific antigen gene (*PSA-TRE*) (U.S. Patent No. 5,648,478), the glandular kallikrein-1 gene (from the human gene, *hKLK2-TRE*), and the probasin gene (*PB-TRE*) (International Patent Application No. PCT/US98/04132). All three of these genes are preferentially expressed in prostate cells and the expression is androgen-inducible. Generally, expression of genes responsive to androgen induction requires the presence of an androgen receptor (AR).

PSA is synthesized exclusively by normal, hyperplastic, and malignant prostatic epithelia; hence, its tissue-specific expression has made it an excellent biomarker for benign prostatic hyperplasia (BPH) and prostatic carcinoma (CaP). Normal serum levels of PSA are typically below 5 ng/ml, with elevated levels indicative of BPH or CaP. Lundwall et al. (1987) *FEBS Lett.* 214: 317; Lundwall (1989) *Biochem. Biophys. Res. Comm.* 161: 1151; and Riegmann et al. (1991) *Molec. Endocrin.* 5: 1921.

The region of the *PSA* gene that is used to provide cell specificity dependent upon androgens, particular in prostate cells, involves approximately 6.0 kilobases. Schuur et al. (1996) *J. Biol. Chem.* 271:7043-7051. An enhancer region of approximately 1.5 kb in humans is located between nt -5322 and nt -3738, relative to the transcription start site of the *PSA* gene. The *PSA* promoter consists of the sequence from about nt -540 to nt +12 relative to the transcription start site. Juxtapositioning of these two genetic elements yield a fully functional, minimal prostate-specific enhancer/promoter (*PSE*) TRE. Other portions of the approximately

6.0 kb region of the *PSA* gene can be used in the present invention, as long as requisite functionality is maintained. In Example 1, adenoviral vector CN796 is described which comprises a composite TRE comprising an HRE and a PSA-TRE, the PSA-TRE comprising a PSA enhancer from -5322 to -3738 fused to a PSA promoter from -541 to +12. This PSA-TRE is derived from adenoviral vector CN706. Rodriguez et al. (1997) *Cancer Research* 57:2559-2563. Accordingly, in one embodiment an adenoviral vector comprises and adenovirus E1A gene under transcriptional control of a composite TRE comprising the cell status-specific TRE, HRE, and a cell type-specific TRE, a PSA-TRE.

The *PSE* and *PSA* TRE used in the present invention are derived from sequences depicted in Figure 4 (SEQ ID NO:3). The enhancer element is nucleotides about 503 to about 2086 of Figure 4 (SEQ ID NO:3). The promoter is nucleotides about 5285 to about 5836 of Figure 4 (SEQ ID NO:3). Accordingly, in some embodiments, the composite TRE comprises an HRE comprising SEQ ID NO:1 and a PSA-TRE comprises nucleotides about 503 to about 2086 of SEQ ID NO:3. In other embodiments, the composite TRE comprises an HRE comprising SEQ ID NO:1 and a PSA-TRE comprises nucleotides about 503 to about 2086 of SEQ ID NO:3 and nucleotides about 5285 to about 5836 of SEQ ID NO:3. As described above, these composite (HRE/PSA) TREs may be operably linked to an adenovirus gene essential for replication, especially an early gene such as E1A or E1B. Example 1 describes such a construct.

In the present invention, replication-competent adenovirus vectors comprising a cell status-specific TRE and a cell type-specific TRE may employ cell type-specific TREs that are preferentially functional in particular tumor cells. Non-limiting examples of tumor cell-specific TREs, and non-limiting examples of respective potential target cells, include TREs from the following genes: α -fetoprotein (*AFP*) (liver cancer), mucin-like glycoprotein DF3 (*MUC1*) (breast carcinoma), carcinoembryonic antigen (*CEA*) (colorectal, gastric, pancreatic, breast, and lung cancers), plasminogen activator urokinase (*uPA*) and its receptor gene (breast, colon, and liver cancers), *HER-2/neu* (*c-erbB2/neu*) (breast, ovarian, stomach, and lung cancers).

Other cell type-specific TREs may be derived from the following exemplary genes (cell type in which the TREs are specifically functional are in parentheses): vascular endothelial growth factor receptor (endothelium), albumin (liver), factor VII (liver), fatty acid synthase (liver), von Willebrand factor (brain endothelium), alpha-actin and myosin heavy chain (both in smooth muscle), synthetase I (small intestine), Na-K-Cl transporter (kidney). Additional cell type-specific TREs are known in the art.

AFP is an oncofetal protein, the expression of which is primarily restricted to developing tissues of endodermal origin (yolk sac, fetal liver, and gut), although the level of its expression varies greatly depending on the tissue and the developmental stage. AFP is of clinical interest because the serum concentration of AFP is elevated in a majority of hepatoma patients, with high levels of AFP found in patients with advanced disease. The serum AFP levels in patients appear to be regulated by AFP expression in hepatocellular carcinoma but not in surrounding normal liver. Thus, the AFP gene appears to be regulated to hepatoma cell-specific expression.

Cell type-specific TREs from the *AFP* gene have been identified. For example, the cloning and characterization of human AFP-specific enhancer activity is described in Watanabe et al. (1987) *J. Biol. Chem.* 262:4812-4818. The entire 5' *AFP* flanking region (containing the promoter, putative silencer, and enhancer elements) is contained within approximately 5 kb upstream from the transcription start site.

The *AFP* enhancer region in human is located between about nt -3954 and about nt -3335, relative to the transcription start site of the *AFP* gene. The human *AFP* promoter encompasses a region from about nt -174 to about nt +29. Juxtapositioning of these two genetic elements yields a fully functional *AFP*-TRE. Ido et al. (1995) describe a 259 bp promoter fragment (nt -230 to nt +29) that is specific for HCC. *Cancer Res.* 55:3105-3109. The *AFP* enhancer contains two regions, denoted A and B, located between nt -3954 and nt -3335 relative to the transcription start site. The promoter region contains typical TATA and CAAT boxes. Preferably, the *AFP*-TRE contains at least one enhancer region. More preferably, the *AFP*-TRE contains both enhancer regions.

Suitable target cells for adenoviral vectors containing *AFP*-TREs are any cell type that allow an *AFP*-TRE to function. Preferred are cells that express, or produce, AFP, including,

but not limited to, tumor cells expressing AFP. Examples of such cells are hepatocellular carcinoma cells, gonadal and other germ cell tumors (especially endodermal sinus tumors), brain tumor cells, ovarian tumor cells, acinar cell carcinoma of the pancreas (Kawamoto et al. (1992) *Hepatogastroenterology* 39:282-286), primary gall bladder tumor (Katsuragi et al. (1989) *Rinsko Hoshasen* 34:371-374), uterine endometrial adenocarcinoma cells (Koyama et al. (1996) *Jpn. J. Cancer Res.* 87:612-617), and any metastases of the foregoing (which can occur in lung, adrenal gland, bone marrow, and/or spleen). In some cases, metastatic disease to the liver from certain pancreatic and stomach cancers produce AFP. Especially preferred are hepatocellular carcinoma cells and any of their metastases. AFP production can be measured using assays standard in the art, such as RIA, ELISA or Western blots (immunoassays) to determine levels of AFP protein production or Northern blots to determine levels of AFP mRNA production. Alternatively, such cells can be identified and/or characterized by their ability to activate transcriptionally an *AFP-TRE* (i.e., allow an *AFP-TRE* to function).

The protein urokinase plasminogen activator (uPA) and its cell surface receptor, urokinase plasminogen activator receptor (uPAR), are expressed in many of the most frequently occurring neoplasia and appear to represent important proteins in cancer metastasis. Both proteins are implicated in breast, colon, prostate, liver, renal, lung and ovarian cancer. Transcriptional regulatory elements that regulate uPA and uPAR transcription have been extensively studied. Riccio et al. (1985) *Nucleic Acids Res.* 13:2759-2771; Cannio et al. (1991) *Nucleic Acids Res.* 19:2303-2308.

CEA is a 180,000-Dalton glycoprotein tumor-associated antigen present on endodermally-derived neoplasia of the gastrointestinal tract, such as colorectal, gastric (stomach) and pancreatic cancer, as well as other adenocarcinomas such as breast and lung cancers. CEA is of clinical interest because circulating CEA can be detected in the great majority of patients with CEA-positive tumors. In lung cancer, about 50% of total cases have circulating CEA, with high concentrations of CEA (greater than 20 ng/ml) often detected in adenocarcinomas. Approximately 50% of patients with gastric carcinoma are serologically positive for CEA.

The 5' upstream flanking sequence of the *CEA* gene has been shown to confer cell-specific activity. The *CEA* promoter region, approximately the first 424 nucleotides upstream of the translational start site in the 5' flanking region of the gene, was shown to confer cell-specific activity when the region provided higher promoter activity in *CEA*-producing cells than in non-producing HeLa cells. Schrewe et al. (1990) *Mol. Cell. Biol.* 10:2738-2748. In addition, cell-specific enhancer regions have been found. WO/95/14100. The entire 5' *CEA* flanking region (containing the promoter, putative silencer, and enhancer elements) appears to be contained within approximately 14.5 kb upstream from the transcription start site. Richards et al. (1995); WO 95/14100. Further characterization of the 5' flanking region of the *CEA* gene by Richards et al. (1995) indicated two upstream regions, -13.6 to -10.7 kb or -6.1 to -4.0 kb, when linked to the multimerized promoter resulted in high-level and selective expression of a reporter construct in *CEA*-producing LoVo and SW1463 cells. Richards et al. (1995) also localized the promoter region to nt -90 and nt +69 relative to the transcriptional start site, with region nt -41 to nt -18 as essential for expression. WO95/14100 describes a series of 5' flanking *CEA* fragments which confer cell-specific activity, such as about nt -299 to about nt +69; about nt -90 to about nt +69; nt -14,500 to nt -10,600; nt -13,600 to nt -10,600, nt -6100 to nt -3800. In addition, cell specific transcription activity is conferred on an operably linked gene by the *CEA* fragment from nt -402 to nt +69, depicted in (SEQ ID NO:6). Any *CEA*-TREs used in the present invention are derived from mammalian cells, including but not limited to, human cells. Thus, any of the *CEA*-TREs may be used in the invention as long as requisite desired functionality is displayed in the adenovirus vector. The cloning and characterization of *CEA* sequences have been described in the literature and are thus made available for practice of this invention and need not be described in detail herein.

The protein product of the *MUC1* gene (known as mucin or MUC1 protein; episialin; polymorphic epithelial mucin or PEM; EMA; DF3 antigen; NPGP; PAS-O; or CA15.3 antigen) is normally expressed mainly at the apical surface of epithelial cells lining the glands or ducts of the stomach, pancreas, lungs, trachea, kidney, uterus, salivary glands, and mammary glands. Zotter et al. (1988) *Cancer Rev.* 11-12: 55-101; and Girling et al. (1989) *Int. J. Cancer* 43: 1072-1076. However, mucin is overexpressed in 75-90% of human breast carcinomas. Kufe et al. (1984) *Hybridoma* 3: 223-232. For reviews, see Hilkens (1988)

Cancer Rev. 11-12: 25-54; and Taylor-Papadimitriou, et al. (1990) *J. Nucl. Med. Allied Sci.* 34: 144-150. Mucin protein expression correlates with the degree of breast tumor differentiation. Lundy et al. (1985) *Breast Cancer Res. Treat.* 5: 269-276. This overexpression appears to be controlled at the transcriptional level.

5 Overexpression of the *MUC1* gene in human breast carcinoma cells MCF-7 and ZR-75-1 appears to be regulated at the transcriptional level. Kufe et al. (1984); Kovarik (1993) *J. Biol. Chem.* 268:9917-9926; and Abe et al. (1990) *J. Cell. Physiol.* 143: 226-231. The regulatory sequences of the *MUC1* gene have been cloned, including the approximately 0.9 kb upstream of the transcription start site which contains a TRE that appears to be involved in
10 cell-specific transcription. Abe et al. (1993) *Proc. Natl. Acad. Sci. USA* 90: 282-286; Kovarik et al. (1993); and Kovarik et al. (1996) *J. Biol. Chem.* 271:18140-18147.

Any *MUC1*-TREs used in the present invention are derived from mammalian cells, including but not limited to, human cells. Preferably, the *MUC1*-TRE is human. In one embodiment, the *MUC1*-TRE may contain the entire 0.9 kb 5' flanking sequence of the *MUC1*
15 gene. In other embodiments, the *MUC1*-TREs comprise the following sequences (relative to the transcription start site of the *MUC1* gene): about nt -725 to about nt +31, nt -743 to about nt +33, nt -750 to about nt +33, and nt -598 to about nt +485 (operably-linked to a promoter).

The *c-erbB2/neu* gene (*HER-2/neu* or *HER*) is a transforming gene that encodes a 185 kD epidermal growth factor receptor-related transmembrane glycoprotein. In humans, the *c-erbB2/neu* protein is expressed during fetal development, however, in adults, the protein is
20 weakly detectable (by immunohistochemistry) in the epithelium of many normal tissues. Amplification and/or over-expression of the *c-erbB2/neu* gene has been associated with many human cancers, including breast, ovarian, uterine, prostate, stomach and lung cancers. The clinical consequences of the *c-erbB2/neu* protein over-expression have been best studied in
25 breast and ovarian cancer. *c-erbB2/neu* protein over-expression occurs in 20 to 40% of intraductal carcinomas of the breast and 30% of ovarian cancers, and is associated with a poor prognosis in subcategories of both diseases. Human, rat and mouse *c-erbB2/neu* TREs have been identified and shown to confer *c-erbB2/neu* expressing cell specific activity. Tal et al. (1987) *Mol. Cell. Biol.* 7:2597-2601; Hudson et al. (1990) *J. Biol. Chem.* 265:4389-4393;

Grooteclaes et al. (1994) *Cancer Res.* 54:4193-4199; Ishii et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:4374-4378; Scott et al. (1994) *J. Biol. Chem.* 269:19848-19858.

The cell type-specific TREs listed above are provided as non-limiting examples of TREs that would function in the instant invention. Additional cell-specific TREs are known in the art, as are methods to identify and test cell specificity of suspected TREs.

Using the adenoviral vectors of the invention

The adenoviral vectors can be used in a variety of forms, including, but not limited to, naked polynucleotide (usually DNA) constructs; polynucleotide constructs complexed with agents to facilitate entry into cells, such as cationic liposomes or other cationic compounds such as polylysine; packaged into infectious adenovirus particles (which may render the adenoviral vector(s) more immunogenic); packaged into other particulate viral forms such as HSV or AAV; complexed with agents (such as PEG) to enhance or dampen an immune response; complexed with agents that facilitate *in vivo* transfection, such as DOTMATM, DOTAPTM, and polyamines. Thus, the invention also provides an adenovirus capable of replicating preferentially in cell status-producing cells. "Replicating preferentially" means that the adenovirus replicates more in cell exhibiting a requisite physiological state than a cell not exhibiting that state. Preferably, the adenovirus replicates at least about 2-fold higher, preferably at least about 5-fold higher, more preferably at least about 10-fold higher, still more preferably at least about 50-fold higher, even more preferably at least about 100-fold higher, still more preferably at least about 400-fold to about 500-fold higher, still more preferably at least about 1000-fold higher, most preferably at least about 1×10^6 higher. Most preferably, the adenovirus replicates solely in cells exhibiting a requisite physiological state (that is, does not replicate or replicates at very low levels in cells not exhibiting the requisite physiological state).

If an adenoviral vector is packaged into an adenovirus, the adenovirus itself may also be selected to further enhance targeting. For example, adenovirus fibers mediate primary contact with cellular receptor(s) aiding in tropism. See, e.g., Amberg et al. (1997) *Virol.* 227:239-244. If a particular subgenus of an adenovirus serotype displayed tropism for a target

cell type and/or reduced affinity for non-target cell types, such subgenus (or subgenera) could be used to further increase cell-specificity of cytotoxicity and/or cytolysis.

The adenoviral vectors may be delivered to the target cell in a variety of ways, including, but not limited to, liposomes, general transfection methods that are well known in the art (such as calcium phosphate precipitation or electroporation), direct injection, and intravenous infusion. The means of delivery will depend in large part on the particular adenoviral vector (including its form) as well as the type and location of the target cells (i.e., whether the cells are *in vitro* or *in vivo*).

If used as a packaged adenovirus, adenovirus vectors may be administered in an appropriate physiologically acceptable carrier at a dose of about 10^4 to about 10^{14} . The multiplicity of infection will generally be in the range of about 0.001 to 100. If administered as a polynucleotide construct (i.e., not packaged as a virus) about 0.01 μg to about 1000 μg of an adenoviral vector can be administered. The adenoviral vector(s) may be administered one or more times, depending upon the intended use and the immune response potential of the host, and may also be administered as multiple, simultaneous injections. If an immune response is undesirable, the immune response may be diminished by employing a variety of immunosuppressants, so as to permit repetitive administration, without a strong immune response. If packaged as another viral form, such as HSV, an amount to be administered is based on standard knowledge about that particular virus (which is readily obtainable from, for example, published literature) and can be determined empirically.

Host cells comprising the adenoviral vectors of the invention

The present invention also provides host cells comprising (i.e., transformed with) the adenoviral vectors described herein. Both prokaryotic and eukaryotic host cells can be used as long as sequences requisite for maintenance in that host, such as appropriate replication origin(s), are present. For convenience, selectable markers are also provided. Prokaryotic host cells include bacterial cells, for example, *E. coli* and mycobacteria. Among eukaryotic host cells are yeast, insect, avian, plant and mammalian. Host systems are known in the art and need not be described in detail herein.

Compositions of the invention

The present invention also provides compositions, including pharmaceutical compositions, containing the adenoviral vectors described herein. Such compositions (especially pharmaceutical compositions) are useful for administration *in vivo*, for example, when measuring the degree of transduction and/or effectiveness of cell killing in an individual. Pharmaceutical compositions, comprised an adenoviral vector of this invention in a pharmaceutically acceptable excipient (generally an effective amount of the adenoviral vector), are suitable for systemic administration to individuals in unit dosage forms, sterile parenteral solutions or suspensions, sterile non-parenteral solutions or oral solutions or suspensions, oil in water or water in oil emulsions and the like. Formulations for parenteral and nonparenteral drug delivery are known in the art and are set forth in *Remington's Pharmaceutical Sciences*, 19th Edition, Mack Publishing (1995). Pharmaceutical compositions also include lyophilized and/or reconstituted forms of the adenoviral vectors (including those packaged as a virus, such as adenovirus) of the invention.

Other compositions are used, and are useful for, detection methods described herein. For these compositions, the adenoviral vector usually is suspended in an appropriate solvent or solution, such as a buffer system. Such solvent systems are well known in the art.

Kits of the invention

The present invention also encompasses kits containing an adenoviral vector(s) of this invention. These kits can be used for diagnostic and/or monitoring purposes, preferably monitoring. Procedures using these kits can be performed by clinical laboratories, experimental laboratories, medical practitioners, or private individuals. Kits embodied by this invention allow someone to detect the presence of cell status-producing cells in a suitable biological sample, such as biopsy specimens.

The kits of the invention comprise an adenoviral vector described herein in suitable packaging. The kit may optionally provide additional components that are useful in the procedure, including, but not limited to, buffers, developing reagents, labels, reacting surfaces, means for detection, control samples, instructions, and interpretive information.

Preparation of the adenovirus vectors of the invention

The adenovirus vectors of this invention can be prepared using recombinant techniques that are standard in the art. Generally, a cell status-specific TRE is inserted 5' to the adenoviral gene of interest, preferably one or more early genes (although late gene(s) may be used). A cell status-specific TRE can be prepared using oligonucleotide synthesis (if the sequence is known) or recombinant methods (such as PCR and/or restriction enzymes). Convenient restriction sites, either in the natural adeno-DNA sequence or introduced by methods such as oligonucleotide directed mutagenesis and PCR, provide an insertion site for a cell status-specific TRE. Accordingly, convenient restriction sites for annealing (i.e., inserting) a cell status-specific TRE can be engineered onto the 5' and 3' ends of a cell status-specific TRE using standard recombinant methods, such as PCR.

Polynucleotides used for making adenoviral vectors of this invention may be obtained using standard methods in the art, such as chemical synthesis, by recombinant methods, and/or by obtaining the desired sequence(s) from biological sources.

Adenoviral vectors are conveniently prepared by employing two plasmids, one plasmid providing for the left hand region of adenovirus and the other plasmid providing for the right hand region, where the two plasmids share at least about 500 nt of middle region for homologous recombination. In this way, each plasmid, as desired, may be independently manipulated, followed by cotransfection in a competent host, providing complementing genes as appropriate, or the appropriate transcription factors for initiation of transcription from a cell status-specific TRE for propagation of the adenovirus. Plasmids are generally introduced into a suitable host cell such as 293 cells using appropriate means of transduction, such as cationic liposomes. Alternatively, *in vitro* ligation of the right and left-hand portions of the adenovirus genome can also be used to construct recombinant adenovirus derivative containing all the replication-essential portions of adenovirus genome. Berkner et al. (1983) *Nucleic Acid Research* 11: 6003-6020; Bridge et al. (1989) *J. Virol.* 63: 631-638.

For convenience, plasmids are available that provide the necessary portions of adenovirus. Plasmid pXC.1 (McKinnon (1982) *Gene* 19:33-42) contains the wild-type left-hand end of Ad5. pBHGI0 (Bett et al. (1994) *Proc. Natl. Acad. Sci USA* 91:8802-8806; Microbix Biosystems Inc., Toronto) provides the right-hand end of Ad5, with a deletion in E3.

The deletion in E3 provides room in the virus to insert a 3 kb cell status-TRE without deleting the endogenous enhancer/promoter. Bett et al. (1994). The gene for E3 is located on the opposite strand from E4 (r-strand). pBHGI1 provides an even larger E3 deletion (an additional 0.3 kb is deleted). Bett et al. (1994).

5 For manipulation of the early genes, the transcription start site of Ad5 E1A is at 498 and the ATG start site of the E1A protein is at 560 in the virus genome. This region can be used for insertion of an cell status-specific TRE. A restriction site may be introduced by employing polymerase chain reaction (PCR), where the primer that is employed may be limited to the Ad5 genome, or may involve a portion of the plasmid carrying the Ad5 genomic
10 DNA. For example, where pBR322 is used, the primers may use the EcoRI site in the pBR322 backbone and the XbaI site at 1339 of Ad5. By carrying out the PCR in two steps, where overlapping primers at the center of the region introduce a 30 sequence change resulting in a unique restriction site, one can provide for insertion of heterologous TRE at that site.

A similar strategy may also be used for insertion of a heterologous TRE to regulate
15 E1B. The E1B promoter of Ad5 consists of a single high-affinity recognition site for SphI and a TATA box. This region extends from 1636 to 1701. By insertion of a heterologous TRE in this region, one can provide for target cell-specific transcription of the E1B gene. By employing the left-hand region modified with a heterologous TRE regulating E1A as the template for introducing a heterologous TRE to regulate E1B, the resulting adenovirus vector
20 will be dependent upon the cell status-specific transcription factors for expression of both E1A and E1B.

Similarly, a cell status-specific TRE can be inserted upstream of the E2 gene to make its expression cell status specific. The E2 early promoter, mapping in Ad5 from 27050-27150, consists of a major and a minor transcription initiation site, the latter accounting for about 5%
25 of the E2 transcripts, two non-canonical TATA boxes, two E2F transcription factor binding sites and an ATF transcription factor binding site. For a detailed review of the E2 promoter architecture see Swaminathan et al., *Curr. Topics in Micro. and Imm.* (1995) 199 (part 3):177-194.

For E4, one must use the right hand portion of the adenovirus genome. The E4
30 transcription start site is predominantly at 35609, the TATA box at 35638 and the first

ATG/CTG of ORF 1 is at 35532. Virtanen et al. (1984) *J. Virol.* 51: 822-831. Using any of the above strategies for the other genes, a cell status-specific TRE may be introduced upstream from the transcription start site. For the construction of mutants in the E4 region, the co-transfection and homologous recombination are performed in W162 cells (Weinberg et al. (1983) *Proc. Natl. Acad. Sci.* 80:5383-5386) which provide E4 proteins *in trans* to complement defects in synthesis of these proteins. Alternatively, these constructs can be produced by *in vitro* ligation.

Methods using the adenovirus vectors of the invention

The adenoviral vectors of the invention can be used for a wide variety of purposes, which will vary with the desired or intended result. Accordingly, the present invention includes methods using the adenoviral vectors described above.

In one embodiment, methods are provided for conferring selective cytotoxicity in target cells (i.e., cells exhibiting a requisite physiological state which allows a cell status-specific TRE to function), generally but not necessarily in an individual (preferably human), comprising contacting the cells with an adenovirus vector described herein, such that the adenovirus vector enters the cell. Cytotoxicity can be measured using standard assays in the art, such as dye exclusion, ³H-thymidine incorporation, and/or lysis.

In another embodiment, methods are provided for propagating an adenovirus specific for mammalian cells which allow a cell status-specific TRE to function. These methods entail combining an adenovirus vector with mammalian cells, whereby said adenovirus is propagated.

The invention further provides methods of suppressing tumor cell growth, generally but not necessarily in an individual (preferably human), comprising contacting a tumor cell with an adenoviral vector of the invention such that the adenoviral vector enters the tumor cell and exhibits selective cytotoxicity for the tumor cell. Tumor cell growth can be assessed by any means known in the art, including, but not limited to, measuring tumor size, determining whether tumor cells are proliferating using a ³H-thymidine incorporation assay, or counting tumor cells.

The invention also includes methods for detecting target cells (i.e., cells which permit or induce a cell status-specific TRE to function) in a biological sample. These methods are particularly useful for monitoring the clinical and/or physiological condition of an individual (i.e., mammal), whether in an experimental or clinical setting. For these methods, cells of a biological sample are contacted with an adenovirus vector, and replication of the adenoviral vector is detected. A suitable biological sample is one in which cells exhibiting a requisite physiological (and/or environmental) state, for example, an aberrant physiological state (such as cells in hypoxic conditions and exhibiting a phenotype characteristic of cells in hypoxic conditions, such as expression of HIF-1) may be or are suspected to be present. Generally, in mammals, a suitable clinical sample is one in which cancerous cells exhibiting a requisite physiological state, such as cells within a solid tumor which are under hypoxic conditions, are suspected to be present. Such cells can be obtained, for example, by needle biopsy or other surgical procedure. Cells to be contacted may be treated to promote assay conditions, such as selective enrichment, and/or solubilization. In these methods, target cells can be detected using *in vitro* assays that detect adenoviral proliferation, which are standard in the art. Examples of such standard assays include, but are not limited to, burst assays (which measure virus yield) and plaque assays (which measure infectious particles per cell). Propagation can also be detected by measuring specific adenoviral DNA replication, which are also standard assays.

The following examples are provided to illustrate but not limit the invention.

EXAMPLES

EXAMPLE 1

Adenovirus vector comprising E1A under transcriptional control of a hypoxia responsive element and a PSA-TRE

General techniques

A human embryonic kidney cell line, 293, efficiently expresses E1A and E1B genes of Ad5 and exhibits a high transfection efficiency with adenovirus DNA. To generate

recombinant adenovirus, 293 cells were co-transfected with one left end Ad5 plasmid and one right end Ad5 plasmid. Homologous recombination generates adenoviruses with the required genetic elements for replication in 293 cells which provide E1A and E1B proteins *in trans* to complement defects in synthesis of these proteins.

5 The plasmids to be combined were co-transfected into 293 cells using cationic liposomes such as Lipofectin (DOTMA:DOPE™, Life Technologies) by combining the two plasmids, then mixing the plasmid DNA solution (10 µg of each plasmid in 500 µl of minimum essential medium (MEM) without serum or other additives) with a four-fold molar excess of liposomes in 200 µl of the same buffer. The DNA-lipid complexes were then placed
10 on the cells and incubated at 37°C, 5% CO₂ for 16 hours. After incubation the medium was changed to MEM with 10% fetal bovine serum and the cells are further incubated at 37°C, 5% CO₂, for 10 days with two changes of medium. At the end of this time the cells and medium were transferred to tubes, freeze-thawed three times, and the lysate was used to infect 293 cells at the proper dilution to detect individual viruses as plaques.

15 Plaques obtained were plaque purified twice, and viruses were characterized for presence of desired sequences by PCR and occasionally by DNA sequencing. For further experimentation, the viruses were purified on a large scale by cesium chloride gradient centrifugation.

Adenovirus vectors in which E1A is under transcriptional control of a cell status-specific TRE

20 An adenovirus vector containing a hypoxia response element (HRE) was generated. CN796, an adenovirus vector in which E1A is under the control of a composite TRE consisting of an HRE and a PSA-TRE, was made by co-transfecting CN515 with pBHGI0. CN515 was constructed by inserting a 67 base pair fragment from HRE eno1 (Jiang et al. (1997) *Cancer Research* 57:5328-5335) (SEQ ID NO:1; Figure 2) into CN65 at the BglIII site. CN65 is a
25 plasmid containing an enhancer and promoter from the human PSA gene, consisting of an enhancer from -5322 to -3738 fused to a PSA promoter from -541 to +12. This is the PSA-TRE contained within plasmid CN706. Rodriguez et al. (1997) *Cancer Res.* 57:2559-2563.

Virus growth in vitro

Growth selectivity of recombinant adenovirus is analyzed in plaque assays in which a single infectious particle produces a visible plaque by multiple rounds of infection and replication. Virus stocks are diluted to equal pfu/ml, then used to infect monolayers of cells for 1 hour. The inoculum is then removed and the cells are overlaid with semisolid agar containing medium and incubated at 37°C for 10 days. Plaques in the monolayer are then counted and titers of infectious virus on the various cells are calculated. The data are normalized to the titer of CN702 (wild type) on 293 cells.

Claims

What is claimed is:

1. An adenovirus vector comprising an adenovirus gene under transcriptional control of a transcriptional regulatory element (TRE) comprising a cell status-specific TRE.
- 5 2. The adenovirus vector of claim 1, wherein the adenovirus gene is essential for viral replication.
3. The adenovirus vector of claim 2, wherein the adenovirus gene is an early gene.
4. The adenovirus vector of claim 2, wherein the adenovirus gene is a late gene.
5. The adenovirus vector of claim 3, wherein the adenovirus early gene is E1A.
- 10 6. The adenovirus vector of claim 3, wherein the adenovirus early gene is E1B.
7. The adenovirus vector of claim 3, wherein the adenovirus early gene is E4.
8. The adenovirus vector of claim 1, wherein the cell status-specific TRE is human.
9. The adenovirus vector of claim 1, wherein the cell status-specific TRE comprises a hypoxia responsive element (HRE).
- 15 10. The adenovirus vector of claim 9, wherein the HRE comprises SEQ ID NO:1.
11. The adenovirus vector of claim 1, wherein the cell status-specific TRE comprises a cell cycle specific element.
12. The adenovirus vector of claim 11, wherein the cell cycle-specific element is from the E2F-1 gene.
- 20 13. The adenovirus vector of claim 1, wherein the cell status-specific TRE comprises a heat-inducible element.

14. The adenovirus vector of claim 1, further comprising a cell type-specific TRE.
15. The adenovirus vector of claim 14, wherein the cell type-specific TRE is prostate cell specific.
- 5 16. The adenovirus vector of claim 15, wherein the prostate cell-specific TRE is a PSA-TRE.
17. The adenovirus vector of claim 1, further comprising a transgene under transcriptional control of a second cell status-specific TRE.
18. An adenovirus vector comprising an adenovirus gene under transcriptional control of a TRE comprising a cell status-specific TRE and a cell-type specific TRE.
- 10 19. The adenovirus vector of claim 18, wherein the adenovirus gene is an early gene.
20. The adenovirus vector of claim 19, wherein the adenovirus early gene is E1A.
21. The adenovirus vector of claim 20, wherein the cell status-specific TRE comprises an HRE and the cell-type specific TRE is a PSA-TRE.
- 15 22. The adenovirus vector of claim 21, wherein the HRE comprises SEQ ID NO:1 and the PSA-TRE comprises nucleotides about 503 to about 2086 of SEQ ID NO:3 and nucleotides about 5285 to about 5836 of SEQ ID NO:3.
23. A composition comprising an adenovirus vector of claim 1.
24. The composition of claim 23, further comprising a pharmaceutically acceptable excipient.
- 20 25. A host cell comprising the adenovirus vector of claim 1.
26. A method of propagating adenovirus specific for cells which allow a cell status-specific TRE to function, said method comprising combining an adenovirus according to claim 1 with the cells, whereby said adenovirus is propagated.

27. A method for conferring selective cytotoxicity on a target cell, said method comprising contacting a cell which allows a cell status-specific TRE to function with an adenovirus vector of claim 1, whereby the vector enters the cell.

- 5 28. A method for suppressing tumor growth comprising introducing the adenovirus vector of claim 1 into a tumor cell which allows a cell status-specific TRE to function, wherein introduction of the adenovirus vector results in suppression of tumor growth.




FIGURE 1

FIGURE 2

ccccgagg cagtgcac gaggtcagg ggggtggt gattcgacgagagaccocg ggggtgcag gccgga

FIGURE 3

99gcccacaa ttacgaatg accactgtg tctaaagca gtggcctaag gaccacctt 61
 gcagaaacgt ggtcccttg tcaagttca ggcagctctt ggttagctt ctgttcttt 121
 cataacctt cccagcgtt gctctggcc agaccagt tggaggagt cgtactgag 181
 ctctagatt ggcaggag gcagatggg aaaaggagt tgtgtgtca gcattggagc 241
 agaggcagca gtggacaata gaggaaatga gtaaatcctt gggagggtc cctagaagt 301
 atgtgtttc ttitttttt tttagacacg gattctcgt tgcctccag gctggtgtg 361
 agtggcatga tcatagctca ctgacgtcc gactctcgg gtccaagcaa tctccacc 421
 tcaagctccc aagtactgag gactacggc acagccacc atgcttgct aatttttga 481
 ttitttag agatgggtt tcaaatgt gatcagct gctcgaact ccgggttca 541
 tggatccac cccgcagct gattacagg attccgggg tagccaccg gcgccagag 601
 ccaattcat gttctgtaaa cgtctgttac ctctgttc cctctctac tggactgtg 661
 gctcttag gccagcaat gggatgggg cacagagcaa gctctccaa cgtttttga 721
 atgagttag gaatgaatga gttcaagcag atgtatacg ttggctgtt gagattttg 781
 caaaaagg acttgacga agcccgacg tcccctcgc cattccag caccctctt 841
 cagcttgcc tctggttag cggataggg ctggtgtcag gattaggata atgcatggg 901
 tgaagcaat tgaggtatga agagtgtgt gatgttggg ctgtggaaat gatgacctg 961
 aaagaagag ggaacagct ctggaatct aagtggagg ctttggggc tacagtttga 1021
 99gtcacgtg cagaacagtg gctctgttct gaactctgac tatagaagg tcsagtggat 1081
 gcggagcgt cggggcgggg cggggcctat gtcccgtt cccacgctt ccagcaggg 1141
 acgcccgggc tggggcgggg gagtacagc gcctcgtt ccatccgac aaagcttgc 1201
 cgcgccccgc cccgcattg gcgcacgc cccgcgccg cgcgccatcc cgcctcgc 1261
 cgcgggtcc ggcggttaa agccaatag aacgcgccg gtgttccc tcacggccg 1321
 ggcagcaat tgtgcggcg ctgcggcgt cgtgtctt tgcggcaaa aagatttgg 1381
 cgcgtaaaag*tgccgggac tttagagga gcggcgccg gggggggag ggtatcgag 1441
 cctgcgcag gctgcgcgc atggggcccg gcgcccgc cgcctgtca ccggggccg 1501
 gcggggcgtg agcgtcatg

FIGURE 4A

aagcttctag tttcttttc cgggtgacat cgtgaaagc actagcatct ctaagcaatg 60
atctgtgaca atattcacag tgtaatgcc tccagggaa ctaactgagc cttgatgtcc 120
agagatTTTT gtgttttttt ctgagactga gtctcgtctt gtgccaggct ggagtgagc 180
ggtgcaacct tggctcaact caagctccgc ctctgggtt cagcccatc tctgtcctca 240
gcctcctgag tagctgggac tacaggcacc cgccaccacg cctggctaata tttttgtat 300
tttttagtag gatgggggtt cactgtgtta gccaggatgg tctcagctc ctgacctcgt 360
gatctgcccc ccttgccctc ccaaatgtct gggatgacag gcgtgagcca ccgcgcctgg 420
ccgatatcca gagatttttt ggggggctcc atcacacaga catgttgac: gtcttcattg 480
ttgactttta gtatccagcc cctctagaaa tctagctgat atagtgtggc tcaaaadctt 540
cagcacaaat cacaccgtta gactatctgg tgtggcccaa accttcaagg gaacaaagg 600
actctaactt ggcaggatat tccaaagcat tagagatgac ctcttgcaaa gaaaaagaaa 660
tggaaaagaa aaagaaaagaa aggaaaaaaa aaaaaaaa gagatgacct ctcaggctct 720
gaggggaaac gcctgaggtc tttgagcaag gtcatcctc tgttgacag tctcctcacc 780
agggtcattg tgacgatcaa atgtgggtcac gtgtatgagg caccagcaca tgctgggtc 840
tggggagtgc cgtgtaagtg tatgcttgca ctgctgaatg cttgggatgt gtcagggtt 900
atcttcagca cttacagatg ctcatctcat cctcacagca tcaatatggg atgggtatta 960
ctggcctcat ttgatggaga aagtggctgt ggctcagaaa ggggggacca ctgaccagg 1020
gacactctgg atgctgggga ctccagagac catgacct caccaactgc agagaaatta 1080
attgtggcct gatgtccctg tcttgaggag ggtggaggtg gaccttcaact aacctctcat 1140
cttgaccctc tcttttaggg ctctttctga cctccacctt ggtactagga ccccatgtta 1200
ttctgtaccc tcttgactct atgaccccca ctgcccactg catccagctg ggtccctcc 1260
tatctctatt cccagctggc cagtgcagtc tcaagtccca cctgtttgtc agtaactctg 1320
aaggggctga catttttactg acttgcaaac aaataagcta actttccaga gtttttgaa 1380
tgctggcaga gtccatgaga ctctgagtc agaggcaagg gcttttactg ctcacagctt 1440
agcagacagc atgaggttca tgttcacatt agtacacctt gcccccccca aatcttgtag 1500
ggtgaccaga cgagctctagg tggatgtctgt gcagaagggg ttttgccac tggtgagaaa 1560
cctgagatta ggaatcctca atcttatact gggacaactt gcaaacctgc tcagcctttg 1620
tctctgtaga agatattatc ttcagatctt tggattgaaa acagacctac tctggaggaa 1680

FIGURE 4B

catattgtat cgattgtcct tgacagtaaa caaatctgtt gtaagagaca ttatctttat 1740
 tatctaggac agtaagcaag cctggatctg agagagatat catcttgcaa ggatgcctgc 1800
 tttaacaaaca tccttgaac aacaatccag aaaaaaaaaa gtgtgtctgt ctttgctcag 1860
 aagacacaca gatacgtgac agaaccatgg agaattgcct cccaacgctg ttcagccaga 1920
 gcttccaccc cttgtctgca ggacagctct aacgttccac cattaaatac gtcttctatc 1980
 acatcctgct tctttatgcc taaccaaggt tctaggctcc gatcgactgt gtctggcagc 2040
 actccactgc caaaccaga ataaggcagc gctcaggatc ccgaaggggc atggctgggg 2100
 atcagaactt ctgggtttga gtgaggagt ggtccacct cttgaatttc aaaggaggaa 2160
 gaggtggat gtgaaggtag tgggggagg aaagtgtcag ttccgaactc ttaggccaat 2220
 gaggaggag actgtaagg tcccagctcc cgaggtactg atgtgggaat ggcctaaga 2280
 tctcatatcc tcaggaaaga ggtgctgaa tctgagggg tagagtctg ggtatattg 2340
 tggcctaagg ctctttggcc cctgaaggca gaggctgaa ccattaggct cagggtttg 2400
 ggtgatagta atgggatctc ttgattcctc aagagtctga ggatcgagg ttgccattc 2460
 ttccatcttg ccacctaate cttactccac ttgagggtat caccagccct cttagctcca 2520
 tgaaggctccc ctgggcaagc acaatctgag catgaaagat gcccagagg ccttgggtgt 2580
 catccactca tcatccagca tcacactctg aggtgtgtgc cagcaccatg acgtcatgtt 2640
 gctgtgacta tccctgcagc gtgctctccc agccacctgc caacogtaga gctgccatc 2700
 ctctctggt gggagtggcc tgcatggtgc caggctgagg cctagtgtca gacagggagc 2760
 ctggaatcat agggatccag gactcaaaag tgctagagaa tggccatag tcaccatcca 2820
 tgaatctca agggcttctg ggtgaggggc acaggaccc gaacttatgg ttcccaagt 2880
 ctattgctct cccaagttag tctccagat acgaggcact gtgccagcat cagccttacc 2940
 tccaccacat cttgtaaaag gactaccag ggcctgatg aacaccatgg tgtgtacagg 3000
 agtaggggtt ggaggcacgg actcctgtga ggtcacagcc aaggagcat catcatgggt 3060
 ggggaggagg caatggacag gcttgagaac ggggatgtgg ttgtatttgg tttctttg 3120
 ttagataaag tgctgggtat aggtatgaga gtggagtatg aagaccagtt aggtaggagg 3180
 atcagattgg agtggtgta gataaagtgc tgggtatagg attgagagt gagtatgaag 3240
 accagttagg atggaggatc agattggagt tgggttagag atggggtaaa attgtgctcc 3300
 ggatgagttt gggattgaca ctgtggaggt ggtttgggat ggcattgctt tgggatgaa 3360

FIGURE 4C

atagatttgt ttgatgttg gctcagacat ccttggggat tgaactgggg atgaagctgg 3420
 gtttgatttt ggaggtagaa gacgtggaag tagctgtcag atttgacagt ggccatgagt 3480
 ttgtttgat ggggaatcaa acaatggggg aagacataag ggttggcttg ttagggttaag 3540
 ttgcgttggg ttgatggggg cggggctgtg tataatgcag ttgatttgt ttgtattaaa 3600
 ttgggttggg tcaggttttg gttgaggatg agttgaggat atgcttgggg acaccggatc 3660
 catgagggtc tcactggagt ggagacaaac ttcctttcca ggatgaatcc agggaaaccc 3720
 taattcacgt gtgaggggag tcaggccact ggctaagtat atccttccac tccagctcta 3780
 agatggtctt aaatttgtat tatctatac cactctgtc tccttcaact tgcttggagt 3840
 ttacctgac actcaactag aaacagggga agattttatc aaattctttt tttttttttt 3900
 ttttttttg gacagagtct cactctgtg ccaggtctg agtgacgtg cgagctctcg 3960
 gctcactgca acctctgcct ccaggttca agtgatctc ctgcctcagc ctctgagtt 4020
 gctgggatta caggcatgca gcaccatgcc cagctaatt ttgtattttt agtagagatg 4080
 ggggttcacc aatgtttgcc aggtctggcct cgaactcctg acctgggtgat ccacctgcct 4140
 cagcctccca aagtgtctgg attacaggcg tcagccaccg cgcccaagcca cttttgtcaa 4200
 attcttgaga cacagctcgg gctggatcaa gtgagctact ctggttttat tgaacagctg 4260
 aaataaccaa ctttttggaa attgatgaaa tcttacggag ttaacagtg aggtaccagg 4320
 gctcttaaga gttcccgatt ctcttctgag actacaaatt gtgattttgc atgccacctt 4380
 aatctttttt tttttttttt taaatcgagg ttctagctc attctatttc ccaggctgga 4440
 gttcaatagc gtgatcacag ctcaactgtg ccttgaaact ctggccttaa gagattctcc 4500
 tgcttcggc tcccaatagc taagactaca gtagtccacc accatatcca gataattttt 4560
 aaattttttt gggggccggg cacagtggct cagcctgta atcccaacac catggggaggc 4620
 tgagatgggt ggaacacgag gtcaggagtt tgagaccagc ctgaccaaca tggtgaaact 4680
 ctgtctctac taaaaaaaa aaaaatagaa aaattagccg ggcgtgggtg cacacggcac 4740
 ctgtaatccc agctactgag gaggtgagg caggagaatc acttgaaccc agaaggcaga 4800
 ggttgcaatg agccgagatt gcgccactgc actccagcct gggtgacaga gtgagactct 4860
 gtctcaaaaa aaaaaaaaa tttttttttt ttgttagaga tggatcttgc tttgtttctc 4920
 tggttggcct tgaactcctg gcttcaagt atcctctac ctggcctcg gaaagtgtg 4980
 ggattacagg cgtgagccac catgactgac ctgtcgtaa tcttgaggta cataaacctg 5040
 gctctaaaag gctaaaaggct aaatatttgt tggagaaggg gcattggatt ttgcattagg 5100

FIGURE 4D

```

atgattctga cctgggaggg caggtcagca ggcattctctg ttgcacagat agagtgtaga 5160
ggtctggaga acaaggagtg gggggttatt ggaattccac attgtttgct gcacgttgga 5220
ttttgaaatg ctagggaact ttgggagact catatttctg ggctagagga tctgtggacc 5280
acaagatctt ttatgatga cagtagcaat gtatctgtgg agctggattc tgggttgga 5340
gtgcaaggaa aagaatgtac taaatgccaa gacatctatt tcaggagcat gaggaataaa 5400
agttctagtt tctggtctca gagtgggtgca gggatcaggg agtctcaca tctctgagt 5460
gctggtgtct tagggcacac tgggtcttgg agtgcaaagg atctaggcac gtgaggcttt 5520
gtatgaagaa tcggggatcg tacccccccc ctgtttctgt ttcatectgg gcatgtctcc 5580
tctgcttttg tcccctagat gaagctctcca tgagctacaa gggcctggtg catccagggt 5640
gatctagtaa ttgcagaaca gcaagtgtcta gctctccctc cccttcaca gctctgggtg 5700
tgaggagggg ttgtccagcc tccagcagca tggggagggc cttggtcagc ctctgggtgc 5760
cagcagggca ggggcggagt cctggggaat gaaggtttta tagggctcct gggggaggct 5820
ccccagcccc aagctt 5836

```

FIGURE 5A

aagcttttta gtgctttaga cagtgaagctg gtctgtctaa cccaagtgaac ctgggctcca	60
tactcagccc cagaagtga ggggaagct ggggtgagcc aaaccaggca agcctaccc	120
cagggctccc agtggcctga gaaccattgg acccaggacc cattacttct agggtaagga	180
aggtacaaac accagatcca accatggctc ggggggacag ctgtcaaatg cctaaaaata	240
tacctggagg aggaagcaggc aaactatcac tgccccaggt tctctgaaca gaaacagagg	300
ggcaacccaa agtccaaatc caggtgagca ggtgcaccaa atgcccaagag atatgacgag	360
gcaagaagtg aaggaaccac ccctgcatca aatgttttgc atgggaagga gaagggggtt	420
gctcatgttc ccaatccagg agaatgcatt tgggatctgc ctctcttctca ctccctgggt	480
agcaagacta agcaaccagg actctggatt tggggaagga cgtttatattg tggagggcag	540
tgtatgacaat ccacgaggg cctaggtgaa gagggcagga aggctcgaga cactggggac	600
tgagtgaaaa ccacacccat gatctgcacc acccatggat gctctctcat tgctacacett	660
tctgttgata tcagatggcc ccattttctg taccttcaca gaaggacaca ggctagggtc	720
tgtgcatggc ctctatcccc ggggccatgt gaggacagca ggtgggaaag atcatgggtc	780
ctcctgggtc ctgcagggcc agaacttca tcaccatac tgacctctca gatgggaatg	840
gcttccctgg ggtcgggcca acggggcctg ggcaggggag aaaggacgtc aggggacagg	900
gagggaaggt catcgagacc cagcctggaa ggttcttgct tctgaccatc caggatttac	960
ttccctgcat ctacctttgg tcattttccc tcagcaatga ccagctctgc ttctgtatct	1020
cagcctccca ccctggacac agcaccaccg tccttgcccc ggctgcatec acccaatacc	1080
ctgataaccc aggaccatt acctctaggg taaggagggt ccaggagaca gaagctgagg	1140
aaaggtctga agaagtaca tctgtcctgg ccagagggga aaaaccatca gatgctgaac	1200
caggagaatg ttgaccagg aaagggaccg aggaaccaag aaaggagta gaccaccagg	1260
gtttgcctga gaggaaggat caaggcccc agggaaagca gggctggctg catgtgcagg	1320
acactgggtg ggcatatgtg tcttagatc tcctgaatt cagtgtccct gccatggcca	1380
gactctctac tcaggcctgg acatgctgaa ataggacaat ggccttgtec tctctcccca	1440
ccattttggca agagacataa aggacattcc aggcacatgcc ttcttgggag gtcacaggtc	1500
tctgtctcac acctcaggga ctgtagttac tgcatcagcc atggtagtg ctgatctcac	1560
ccagctctgc caggcccttc cactctccac ttgttgacca tgtccaggac caccctcag	1620
atctctgagcc tgcaaatacc ccttctgctg gtgggtggat tcagtaaaaca gtgagctct	1680

FIGURE 5B

atccagcccc	cagagccacc	tctgtcacct	tctgtctggg	catcatccca	ccttcacaag	1740
cactaaagag	catggggaga	cctggctagc	tgggtttctg	catcacaaag	aaaataatcc	1800
cccaggtctg	gattccccag	gctctgtatg	tggagctgac	agacctgagg	ccaggagata	1860
gcagagggtca	gccctagggg	gggtgggtca	tcaccccagg	ggacaggggt	gcaccagcct	1920
tgctactgaa	agggcctccc	caggacagcg	ccatcagccc	tgccctgagag	ctttgctaaa	1980
cagcagtcag	aggaggccat	ggcagtggtc	gagctcctgc	tcaggcccc	aacagaccag	2040
accaacagca	caatgcagtc	cttcccacac	gtcacaggtc	accaaggga	aactgaggtg	2100
ctacctaac	ttagagccat	caggggagat	aacagcccaa	tttcccaaac	aggccagttt	2160
caatcccatg	acaatgacct	ctctgctctc	attcttccca	aaataggacg	ctgattctcc	2220
cccaccatgg	atttctccct	tgtcccggga	gccttttctg	ccccctatga	tctgggacct	2280
cctgacacac	acctctctct	tgggtacata	tcagggtccc	tcactgtcaa	gcagtcacaga	2340
aaggacagaa	ccttgagcag	cgcccatctc	agcttcaccc	ttcctctctc	acaggggttca	2400
gggcaaaagaa	taaatggcag	aggccagtga	gcccaagatg	ggtgacaggc	agtgaccacg	2460
gggcagatgc	ctggagcagg	agctggcggg	gccacaggga	gaagggtgatg	cagggaaggga	2520
aaccagaaa	tgggcaggaa	aggaggacac	aggctctgtg	gggctgcagc	ccaggggttg	2580
actatgagtg	tgaagccatc	tcagcaagta	aggccaggtc	ccatgaacaa	gagtgaggagc	2640
acgtggcttc	ctgctctgta	tatggggtgg	gggattccat	gccccataga	accagatggc	2700
cggggttcag	atggagaagg	agcaggacag	gggatcccca	ggataggagg	accccagtg	2760
cccccccag	gcaggtgact	gatgaatggg	catgcagggt	ectcctgggc	tgggctctcc	2820
ctttgtccct	caggattcct	tgaaggaa	tcgggaagcc	gaccacatct	acctggtggg	2880
ttctggggag	tccatgtaaa	gccaggagct	tgtgttgcta	ggaggggtca	tggcatgtgc	2940
tgggggcacc	aaagagagaa	acctgagggc	aggcaggacc	tggctgagg	aggcatggga	3000
gccagatgg	ggagatggat	gtcaggaaag	gctgccccat	cagggaagggt	gatagcaatg	3060
gggggtctgt	gggagtgggc	acgtgggatt	ccctgggctc	tgccaagttc	cctcccatag	3120
tcacaacctg	gggaactgc	ccatgaaggg	ggccttttgc	ccagccagat	gctgctggtt	3180
ctgccatctc	actaccctct	ctgctccagc	cactctgggt	ctttctccag	atgccctgga	3240
cagccctggc	ctgggcctgt	ccctgagag	gtgtggggag	aagctgagtc	tctggggaca	3300
ctctcatcag	agtcgaaag	gcacatcagg	aaacatccct	ggtctccagg	actaggcaat	3360

FIGURE 5C

gaggaagg	ccccagctcc	tccctttgcc	actgagaggg	tcgacccctgg	gtggccacag	3420
tgacttctg	gtctgtccca	gtcaccctga	aaccacaaca	aaaccccgag	cccagacccct	3480
gcaggtagaa	tacatgtggg	gacagtctgt	accaggggga	agccagttct	ctcttctctag	3540
gagaccgggc	ctcaggctg	tgcccggggc	aggcgggggc	agcacgtgcc	tgctcctttag	3600
aactcgggac	cttaagggtc	tctgctctgt	gaggcacagc	aaggtacctt	ctgtccagag	3660
atgaaagcag	ctcctgcccc	tcctctgacc	tcttctctct	tcccaaatct	caaccaacaa	3720
ataggtgttt	caaatctcat	catcaaatct	tcattccatcc	acatgagaaa	gcttaaaacc	3780
caatggattg	acaacatcaa	gagttggaac	aagtggacat	ggagatgtta	cttgtggaaa	3840
tttagatgtg	ttcagctatc	gggcaggaga	atctgtgtca	aattccagca	tggttcagaa	3900
gaatcaaaaa	gtgtcacagt	ccaaatgtgc	aacagtgcag	gggataaaac	tgtggtgcat	3960
tcaaactgag	ggatattttg	gaacatgaga	aaggaaagga	ttgctgtctg	acagaacatg	4020
gatgatctca	cacatagagt	tgaagaaag	gagtcaatcg	cagaatagaa	aatgatcact	4080
aattccacct	ctataaagtt	tccaagagga	aaacccaatt	ctgctgctag	agatcagaat	4140
ggaggtgacc	tgctgccttg	aatggctgtg	agggtcacgg	gagtgctact	tagtgacggc	4200
aatgtgccgt	atcttaaatct	gggcagggct	ttcatgagca	cataggaatg	cagacattac	4260
tgctgtgttc	attttacttc	accggaaaaag	aagaataaaa	tcagccgggc	gcggtggctc	4320
acgcctgtaa	tcccagcact	ttagaaggct	gaggtgggca	gattacttga	ggtcaggagt	4380
tcaagaccac	cctgcgccaat	atggtgaaac	cccggtctcta	ctaaaaatac	aaaaattagc	4440
tgggcatggt	ggtgcgcgcc	tgtaatccca	gctactcggg	aggctgaggc	tggaacaattg	4500
cttggaacca	ggaagcagag	gttgccagtga	gccaaagtgt	tgccactgca	ctccagcttg	4560
ggcaacagag	ccagactctg	taaaaaaaaa	aaaaaaaaaa	aaaaaaagaa	agaaagaaaa	4620
agaaagaaaa	gtataaaatc	tctttgggtt	aacaaaaaaa	gatccacaaa	acaaacacca	4680
gctcttatca	aacttacaca	actctgccag	agaacaggga	acacaaatac	tcattaactc	4740
acttttgtgg	caataaaacc	ttcatgtcaa	aaggagacca	ggacacaatg	aggaagttaa	4800
actgcagacc	ctacttgggt	gcagagaggg	aaaatccaca	aataaaaacat	taccagaagg	4860
agctaagatt	tactgcattg	agttcattec	ccaggtatgc	aaggtgattt	taaaccttga	4920
aaatcaatca	ttgcctttac	tacatagaca	gattagctag	aaaaaaatta	caactagcag	4980
aacagaagca	attttgcctt	cctaaaaatc	cacatcatat	catcatgatg	gagacagtg	5040
agacgccaat	gacaataaaa	agagggacct	ccgtcaccgc	gtaaaacatgt	ccacacagct	5100

FIGURE 5D

ccagcaagca cccgtcttcc cagtgaatca ctgtaacctc cccctttaatc agccccaggg	5160
aaggctgcct gcgatggcca cacaggctcc aacctgtggg cctcaacctc ccgcagaggg	5220
tctcctttgg ccccccatg gggagagcat gaggacaggg cagagccctc tgatgccac	5280
acatggcagg agctgacgac agagccatgg gggctggaga gcagagctgc tggggtcaga	5340
gcttctctgag gacacccagg cctaaggaa ggcagctccc tggatggggg caaccaggct	5400
ccgggctcca accctagagc ccgcatggga ggagccagca ctctaggcct ttcttagggt	5460
gactctgagg ggacctgac acgacaggat cgctgaatgc acccgagatg aaggggccac	5520
acggggaccc tgctctcgtg gcagatcagg agagagtggg acaccatgcc agggcccat	5580
ggcatggctg cgactgaccc agggcaactcc cctgcagtca tcagcctcgg taagtacat	5640
gaccaagccc aggaccaatg tggaaaggaa gaaacagcat cccctttagt gatggaaccc	5700
aaggctcagt caaagagagg ccatgagcag ttagggaagg tgggtccaacc tacagcacia	5760
accatcatct atcataagta gaagccctgc tccatgaccc ctgcatttaa ataacggtt	5820
gttaaatgag tcaaattccc tcaccatgag agctcacctg tgtgtaggcc catcacacac	5880
acaacacac acacacacac acacacacac acacacacac acagggaaag tgcaggatcc	5940
tgacacagca caggcaggct tcacagcgag agcaaacagc gtgaatgacc catgcagtgc	6000
cctgggcccc atcagctcag agacctgtg agggctgaga tggggctagg caggggagag	6060
acttagagag ggtggggcct ccaggggagg ggcctgcagg agctgggtac tggcctccag	6120
ggggggggct gcaggggagct gggtaactgc ctccaggagg ggggctgcag ggagctgggt	6180
actgccctcc agggaggggg ctgcaggagg ctgggtactg cctccagggg agggggctgc	6240
agggagctgg gtactgcctt ccaggggagg agggagctgc ttcccaacag agagcacatc	6300
ttctgcagc agctgcacag acacaggagc ccccatgact gccctgggcc aggggttgg	6360
ttccaaattt cgtgccccat tgggtgggac ggaggttgac cgtgacatcc aaggggcac	6420
tgtgattcca aactaaact actgtgccta caaaatagga aataacccta cttttctac	6480
tatctcaaat tccctaagca caagctagca ccccttaaat caggaaagtc agtcactcct	6540
ggggctctcc catgccccca gtctgacttg caggtgcaca ggggtgctga catctgtcct	6600
tgctctcctt cttggctcaa ctgccgcccc tccctggggg gactgatggt caggacaagg	6660
gactctagag ctggcccat gattgacagg aaggcaggac ttggcctcca ttctgaagac	6720
taggggtgtc aagagagctg ggcacccac agagctgcac aagatgacgc ggacagaggg	6780

FIGURE 5E

tgacacaggg ctcagggctt cagacgggtc gggaggctca gctgagagtt caggggacaga	6840
cctgaggagc ctcagtggga aaagaagcac tgaagtggga agttctggaa tgttctggac	6900
aagcctgagt gctctaagga aatgctccca cccgatgta gcctgcagca ctggacggtc	6960
tgtgtacctc cccgctgccc atcctctcac agccccgcc tctagggaca caactcctgc	7020
cctaaccatgc atctttcctg tctcattcca cacaaaaggc cctctggggt cctgtttctg	7080
cattgcaagg agtggaggtc acgttccac agaccacca gcaacagggt cctatggagg	7140
tgcggtcagg aggatcacac gtcccccat gcccaggga ctgactctgg gggatgtgga	7200
ttggcctgga ggccactggt cccctctgtc cctgaggga atctgcacc tgagggtctgc	7260
cacatccctc ctgattcttt cagctgaggc cctttctga aatcccagg aggactcaac	7320
ccccactggg aaaggcccag tgtggacggt tccacagcag ccagctaaag gccttggac	7380
acagatcctg agtgagagaa cctttaggga cacagggtga cggccatgtc cccagtgcgc	7440
acacagagca ggggcatctg gacccctgagt gtgtagctcc cgcgactgaa cccagccctt	7500
ccccatgac gtgacccctg gggtagctcc aggtctccag tccatgccac caaaatctcc	7560
agattgaggg tcctcccttg agtccctgat gcctgtccag gagctgcccc ctgagcaaat	7620
ctagagtga gagggtggg attgtggcag taaaagcagc cacatttctc tcaggaagga	7680
aaggaggagc atgagctcca ggaaggcgga tggcgctctc tagtgggcgc ctctgttaa	7740
tgagcaaaaa ggggccagga gagttagag atcagggctg gccttggaat aaggctcaga	7800
tggagaggac tgagggtcaa agagggggct gaagtagggg agtggtcggg agagatggga	7860
ggagcaggtg aggggaagcc ccaggaggc cgggggaggg tacagcagag ctctccactc	7920
ctcagcattg acatttgggg tggctgtgct agtggggttc tgtaagttgt aggggttca	7980
gcaccatctg gggactctac ccaactaaat ccagcaggac tccctcccca agctctaaca	8040
accaacaatg tctccagact ttccaaatgt cccctggaga gcaaaattgc tcttggcaga	8100
atcactgatc tacgtcagtc tctaaaagt actcatcagc gaaatccttc acctcttggg	8160
agaagaatca caagtgtgag aggggtagaa actgcagact tcaaaatctt tccaaaagag	8220
ttttacttaa tcagcagttt gatgtcccag gagaagatac atttagagtg tttagagttg	8280
atgccacatg gctgcctgta cctcacagca ggagcagagt gggttttcca agggcctgta	8340
accacaactg gaatgacact cactgggtta cattacaaag tggaatgtgg ggaattctgt	8400
agactttggg aagggaaatg tatgacgtga gccacagcgc taaggcagtg gacagtccac	8460
tttgaggctc tcaccatcta ggagacatct cagccatgaa catagccaca tctgtcatta	8520

FIGURE 5F

gaaaacatgt	tttattaaga	ggaaaaatct	aggctagaag	tgccttatgc	tcttttttct	8580
ctttatgttc	aaattcatat	acttttagat	cattccttaa	agaagaatct	atccccctaa	8640
gtaaatgtta	tcactgactg	gatagtgttg	gtgtctcact	ccccaccctc	gtgtggtgac	8700
agtgcoctgc	tccccagcc	ctgggccctc	tctgattcct	gagagctttg	ggtgctcctt	8760
cattaggagg	aagagaggaa	gggtgttttt	aatattctca	ccattcacc	atccacctct	8820
tagacactgg	gaagaatcag	ttgccactc	ttggatttga	tcctcgaatt	aatgacctct	8880
atttctgtcc	ctgtccatt	tcaacaatgt	gacaggccta	agaggtagct	tctccatgtg	8940
atttttgagg	agaaggttct	caagataagt	tttctcacac	ctctttgaat	tacctccacc	9000
tgtgtcccca	tcaccattac	cagcagcatt	tggacccttt	tctgttagt	cagatgcttt	9060
ccacctcttg	aggggtgtata	ctgtatgtct	tctacacagg	aatatgcaga	ggaatatagaa	9120
aaagggaat	cgcattacta	ttcagagaga	agaagacctt	tatgtgaatg	aatgagagtc	9180
taaaactcta	agagagccca	tataaaatta	ttaccagtgc	taaaactaca	aaagttacac	9240
taacagtaaa	ctagaataat	aaaacatgca	tcacagttgc	tggtaaagct	aaatcagata	9300
tttttttctt	agaaaaagca	ttccatgtgt	gttgacagtga	tgacaggagt	gcccttcagt	9360
caatatgctg	cctgtaattt	ttgttcctcg	gcagaatgta	ttgtcttttc	tccttttaaa	9420
tcttaaatgc	aaaactaaag	gcagctcctg	ggccccctcc	ccaaagtccg	ctgcctgcaa	9480
ccagccccac	gaagagcaga	ggcctgagct	tccttggtca	aaataggggg	ctaggaggct	9540
taacctgtct	cgataaagct	gtgttcccg	aatgtcgcct	ctgttcccg	gggcaccagc	9600
ctggagggtg	gtgagcctca	ctgttgccct	gatgcttacc	ttgtgccctc	acaccagtgg	9660
tcactggaac	cttgaacatt	tggctgtcgc	ccggtatctgc	agatgtcaag	aactctgga	9720
agtcaaatat	ctgccacctt	ctccagggca	gatacctgtg	aacatccaaa	accatgccac	9780
agaacctctg	ctgggggtcta	caacacatat	ggactgtgag	caccaagtcc	agccctgaat	9840
ctgtgaccac	ctgccaaagt	gcccttaact	gggatccacc	aatcactgca	catggcaggc	9900
agcgaggctt	ggaggtgctt	cgccacaagg	cagccccaat	ttgctgggag	tttcttgga	9960
octggtagtg	gtgaggagcc	ttggggacct	caggattact	ccccttaagc	atagtgggga	10020
cccttctgca	tccccagcag	gtgccccgct	cttcagagcc	tctctctctg	aggtttaccc	10080
agacctctgc	accaatgaga	ccatgctgaa	gcctcagaga	gagagatgga	gctttgacca	10140
ggagccgctc	tctcttgagg	gccagggcag	ggaagcagg	aggcaagacc	aggagtggga	10200

FIGURE 5G

```

acaccagtgt ctaagcccct gatgagaaca ggggtgtctc tcccatatgc ccataccagg 10260
cctgtgaaca gaatcctcct tctgcagtga caatgtctga gaggacgaca tgtttccag 10320
cctaactgtc agccatgcc atctaccac tgctactgc aggacagcac caaccaggga 10380
gctgggaagc tgggagaaga catggaatac ccattgcttc tcaccttctt ccagtcagat 10440
gggcaccatt tatgcctagg acaccacact gccggcccca ggtctttaag agttaggtca 10500
cctaggtgcc tctgggaggc cgaggcagga gaattgcttg aaccgggag gcagaggttg 10560
cagtgcgcgc agatcacacc actgcactcc agcctgggtg acagaatgag actctgtctc 10620
aaaaaaaaa agaaagatag catcagtggc taccaagggc taggggcagg ggaaggtgga 10680
gagttaatga ttaatatgat gaagtttcta tgtgagatga tgaaaaatgt ctgaaaaaaa 10740
aaatatagtg gtgaggatgt agaatttgt gaataaatt aacggcattt aattgtacac 10800
ttaacatgat taattgggca tattttatct tatgtatttg actacatcca agaaacactg 10860
ggagagggaa agcccaccat gtaaaataca cccaccctaa tcagatagtc ctcatgttac 10920
ccaggtacag gccctcatg acctgcacag gaataactaa ggatttaagg acatgaggct 10980
tcccagcaa ctgcaggtgc acaacataaa tgtatctgca aacagactga gagtaaagct 11040
gggggcacaa acctcagcac tgccaggaca cacacccttc tcgtggattc tgactttatc 11100
tgaccgggcc cactgtccag atcttgttgt gggattggga caagggaggt cataaagcct 11160
gtcccagggt cactctgtgt gagcacaga gacctcccca cccccccacc gttaggtctc 11220
cacacataga tctgaccatt aggcatttgt aggggaactc tagcgcgggc tcaggggatca 11280
caccagagaa tcaggtacag agagggaagc ggggctcgag gagctgatgg atgacacaga 11340
gcagggttcc tgcatccac aggtccagct caccctgggt taggtgcccc atccccctga 11400
tccaggcatc cctgacacag ctccctccc gagctcctc ccaggtgaca catcagggtc 11460
cctcaactca gctgtccaga gagggcagca cttggacag cgccaccccc acttcaactc 11520
tcctccctca cagggtccag ggctcagggc tcaagtctca gaacaaatgg caggaggcag 11580
tgagcccaga gatggtgaca gggcaatgat ccaggggcag ctgcctgaaa cgggagcagg 11640
tgaagccaca gatgggagaa gatggttcag gaagaaaaat ccagggaatgg gcaggagagg 11700
agaggaggac acaggctctg tggggctgca gccaggatg ggactaagtg tgaagacatc 11760
tcagcagggt aggcagggtc ccattgaacag agaagcagct cccacctccc ctgatgcacg 11820
gacacacaga gtgtgtggtg ctgtgcccc agagtgcggc tctcctgttc tggctccccg 11880
ggagtgaaga gtgaggttga cttgtccctg ctctctctg ctaccccaac attcaacttc 11940

```

FIGURE 5H

tcctcatgcc cctctctctc aaatatgatt tggatctatg tccccgccca aatctcatgt	12000
caaaattgtaa accocaaatg tggagggtggg gccttgtgag aagtgattgg ataatgcggg	12060
tggattttct gctttgatgc tgtttctgtg atagagatct cacatgatct ggttgtttaa	12120
aagtggttag cacctctccc ctctctctct ctctctctta ctcatgctct gccatgtaa	12180
acgttctctgt ttccccttca ccgtccagaa tgattgtaa ttttctgagg cctcccagg	12240
agcagaagcc actatgcttc ctgtacaact gcagaatgat gagcgaatta aacctcttt	12300
ctttataaat taccagctct caggtatttc ttatagcaa tgcgaggaca gactaatata	12360
atcttctact cccagatccc cgcacacgct tagcccaga catcactgcc cctgggagca	12420
tgcacagcgc agcctctctc gcacaaaagc aaagtcacaa aaggtgacaa aaatctgcat	12480
ttggggacat ctgattgtga aagagggagg acagtacact ttagccaca gagactgggg	12540
ctcacccagc tgaaccttgg tagcactttg gcataacatg tgcagtaccc gtgtcaatg	12600
tctagagatc agtgttgagt aaaacagcct ggtctggggc cgctgctgtc cccacttccc	12660
tctgtctcac cagagggcgg cagagttctc ccacccctg agcctcccga ggggtgctg	12720
acctcccctc gccgggccca cagcccagca gggccaacc tcaccgggt cacctggcc	12780
cacgtctctc tcgcctctcg agctctcac acggaactct tcagctctc cctgcagcct	12840
atcgccgcc cacttgaggc ttgtcggcgc ccaacttgag gctgtcggc tgcctctgc	12900
aggcagctcc tgtcccctac accccctctc tcocgggct cagctgaaag ggcgtctccc	12960
agggcagctc cctgtgatct ccaggacagc tcagttcttc acaggtccc agcccctta	13020
tgtctcacc tcacagccct gtcattacca ttaactctc agtccatga agttcactga	13080
gcgcctgtct cccggttaca ggaactctc gtgacagga ccacgtctgt cctgtctct	13140
gtggaatccc agggcccagc ccagtgcctg acacggaaac gatgtccat aaatactggt	13200
taaaatgtgt ggagatctct aaaaagaagc atatcacctc cgtgtggccc ccagcagtc	13260
gagttctgtc catgtgtgga caggggcact ggcaccagca tgggagggag ccagcaagt	13320
cccgcggctg ccccaggaat gaggcotcaa ccccagagc tcagaaggg aggacagag	13380
cctgcagggg atagatctct cggcctgacc ctgcagccta atccagagtt cagggtcagc	13440
tcacaccacg tgcaccttg tcagcatccc tagggcagtt ccagacaagg ccggaggtct	13500
cctcttgccc tcacgggggt gacattgcac acagacatca ctacggaaac ggattcccct	13560
ggacaggaac ctggctttgc taagggaagt gaggtggagc ctggtttcca tcccttgcct	13620

FIGURE 5I

caacagacccc	ttctgatctc	tcccacatac	ctgctctgtt	cctttctggg	tcttatgagg	13680
accctgttct	gccaggggtc	cctgtgcaac	tccagactcc	ctcctggtag	caccatgggg	13740
aaggtggggg	gatcacagga	cagtcagcct	cgcagagaca	gagaccaccc	aggactgtca	13800
gggagaacat	ggacaggccc	tgagcccgag	ctcagccaac	agacacggag	agggaggggtc	13860
ccctcggagc	cttccccaag	gacagcagag	cccagagtca	cccacctccc	tccaccacag	13920
tcctctcttt	ccaggacaca	caagacacct	ccccctccac	atgcaggatc	tggggactcc	13980
tgagacctct	gggcctgggt	ctccatccct	gggtcagtag	cgggggttgt	ggtactggag	14040
acagaggggt	ggtccctccc	cagccaccac	ccagttagcc	ttttctagc	ccccagagcc	14100
acctctgtca	ccttctctgt	gggcatcacc	ccacctcccc	agagccctgg	agagcatggg	14160
gagacccggg	accctgctgg	gtttctctgt	cacaaaggaa	aataatcccc	ctgggtgtgac	14220
agacccaagg	acagaacaca	gcagagggtca	gcactggggg	agacagggtg	tcctcccagg	14280
ggatgggggt	ccatccacct	tgccgaaaag	atttgtctga	ggaactgaaa	atagaagggg	14340
aaaaagagga	gggacaaaag	aggcagaaat	gagaggggag	gggacagagg	acacctgaat	14400
aaagaccaca	cccatgaccc	acgtgatgct	gagaagtact	cctgccctag	gaagagactc	14460
agggcagagg	gaggaaggac	agcagaccag	acagtcacag	cagccttgac	aaaacgttcc	14520
tggaactcaa	gctcttctcc	acagaggagg	acagagcaga	cagcagagac	catggagtct	14580
ccctcggccc	ctccccacag	atggtgcacc	ccctggcaga	ggctcctgct	cacaggtgaa	14640
gggaggacaa	cctggggagag	gggtgggagga	gggagctggg	gtctcctggg	taggacaggg	14700
ctgtgagagc	gacagagggc	tcctgttgga	gcctgaatag	ggaagaggac	atcagagagg	14760
gacaggagtc	acaccagaaa	aatcaaatg	aactggaatt	ggaaaagggc	aggaaaacct	14820
caagagttct	attttctctag	ttaattgtca	ctggccacta	cgtttttaaa	aatcataata	14880
actgcatacg	atgacacttt	aaataaaaac	ataaccaggg	catgaaacac	tgtcctcacc	14940
cgctaccgcg	ggacattgga	aaataagccc	caggctgtgg	agggccctgg	gaacctcat	15000
gaactcatcc	acaggaatct	gcagcctgtc	ccaggccactg	gggtgcaacc	aagatc	15056

FIGURE 6A

gaattcagaa ataggggaag gttgaggaag gacactgaac tcaaagggga tacagtatt 60
 gggtttatttg tcttctcttc acaacattgg tgctggagga attcccaccc tgaggttatg 120
 aagatgtctg aacacccaac acatagcact ggagatatga gctcgacaag agtttctcag 180
 ccacagagat tcacagccta gggcaggagg acactgtacg ccaggcagaa tgacatggga 240
 attgcctcga cgattggcct gaagaagcaa ggactgtggg aggtgggctt ttagtaaca 300
 agaggcgagc gtgaactctg attcccatgg ggaatgtga tggctcgtt acaaaatttt 360
 caagctggca ggaataaaaa cccattacgg tgaggacctg tggaggcggt ctgccccaac 420
 tgataaagga aatagccagg tgggggcctt tccattgta ggggggacat atctggcaat 480
 agaagccttt gagacccttt aggggtacaag tactgaggca gcaataaaaa tgaatctta 540
 ttttcaact ttatactgca tgggtgtgaa gatataattg tttctgtaca gggggtgag 600
 gaaaggaggg gaggaggaaa gttcctgcag gtctggtttg gtctgtgat ccagggggtc 660
 ttggaactat ttaataaaa ttaataaaa acaagcagct gtttaaat aaattaaatt 720
 aaattaaatt ttactttatt ttatottaag ttctgggcta catgtgcagg acgtgcagct 780
 ttgttacata ggtaaaacgtg tgccatgggt gtttgcgtga cctatcaacc catcacctag 840
 gtattaagcc cagcatgcat tagctgtttt tctgacgct ctccctctcc ctgactccca 900
 caacaggccc cagtggtgtg tgttccctc cctgtgtcca tgtgttctca ttgttcagct 960
 cccacttata agtgagaaca tgtggtgttt ggttttctgt ttctgtgtta gtttgcag 1020
 gataatggct tccactcca tccatgttcc tgcaaggagc gtgatcttat tctttttat 1080
 ggttgcatag aaattgtttt tacaatcca attgatattg tatttaatta caagttaatc 1140
 taattagcat actagaagag attacagaag atattaggtg cattgaatga ggaatatat 1200
 aaaataggac gaagtgaaa tattaggtag gaaaagtata atagttgaaa gaagtaaaaa 1260
 aaaatatgca tgagtagcag aatgtaaaag aggtgaagaa cgtaatagtg actttttaga 1320
 ccagattgaa ggacagagac agaaaaattt taaggaaatg ctaaacatg tgagtgttag 1380
 aagtacagtc aataacatta agcctcagg agggagaaa aataggaaa gaggaaatat 1440
 gtgaataaat agtagagaca tgtttgatgg attttaaat atttgaaaga cctcacatca 1500
 aggatccat accgtgccat tgaaggagaa gatggaaa ccaagaagcc agatgaaagt 1560
 tagaaatatt attggcaaa cttaaatggt aaaagtccta gagagaaagg atggcagaaa 1620
 tattggcggg aaagaatgca gaacctagaa tataaatcca tcccaacagt ttgtagtgt 1680

FIGURE 6B

gcagctgtag ccttttctag ataatacact attgtcatac atcgcttaag cgagtgtaaa 1740
atggtctcct caacttattt atttatatat ttatttagtt ttgagatgga gccctgctct 1800
gtctcctagg ctggagtgca atagtgcgat accactcact gcaacctctg cctcctctgt 1860
tcaagtattt ttcttacctc agcctcccca gtactgggga ttacaggtgc gtgccaccac 1920
accgggctaa tttttgtatt tttttagag acgggggttt gccatgttg ccagctggt 1980
cttgaactcc tgacatcagg tgatccacct gccctggcct cctaaagtgc tgggattaca 2040
ggcatgagcc accgtgcccc accactttat ttatttttta tttttattt taaatttcag 2100
cttctatttg aaatacaggg ggcacatata taggattgtt acatgggtat attgaactca 2160
ggtagtgatc atactaccac acaggtagggt ttcaaccaca ctccccctct tttctcccc 2220
attctagtag tgtgcagtgt ctattgttct catgtttatg tctatgtgtg ctccagggtt 2280
agctccacc ttgaagttag aacgtgtggt atttgatttt ctgtccctgt gttattccac 2340
ttaggattat ggcttccagc tccattcata ttgtctgaaa ggatagatt catttttcat 2400
ggccatgcag tattccatat tgcgtataga tcacattttc tttcttttt ttttttga 2460
cggagtcttg ctttctctgc taggctggag tgcagttagca cgatctcggc tcactgcaag 2520
cttcacctcc ggggttccag tcattcttct gtctcagctt cccaagtgc tgggactaca 2580
gggcgccgcc accacgtccg gctaattttt ttgtgtgttt ttagtagaga tgggggttc 2640
actgtgttag ccaggtggt ctgactctcc tgaccttggt gtccacctgc ctcggtctcc 2700
caaatgtctg ggtattacagg ggtgagccac tgcgcccgcc ccatatatac cacttttct 2760
ttaaccaatc caccattgat gggcaactag gtatattcca tggattccac agttttgcta 2820
ttgtgtcag tgtggcagta gacatatgaa tgaatgtgtc tttttggtat aatgatttgc 2880
attcctttgg gtatacagtc attaatagga gtgtgtgtgt gaacgtgtgc tctgtttaaa 2940
attctttgag aattttccaa actgtttgccc atagagagca aactaattta catttccagc 3000
aacagtatat aagcattccc ttttctccac agctttgtca tcatggtttt ttttttctt 3060
tattttaaaa aagaatatgt tgtgttttcc ccagggtaca tgtgcaggat gtgcagggtt 3120
gttcatatagg tagtaaacgt gagccatggt ggtttgctgc acctgtcaac ccattacctg 3180
ggatgaagc cctgcctgca ttactctttt tccctaatgc tctcactact gcccccacct 3240
cacctgaca gggcaaacag acaacctaca gaatgggagg aaatttttgc aatctattca 3300
cttgacaaag gtcaagaata tccagaatct acaaggaaat taagcaaat tttacttttt 3360

FIGURE 6C

aataatagcc actctgactg gcgtgaaatg gtatctcatt gtggttttca ttggaatttc 3420
tctgatgac agtgacgatg agcatttttt catatttggt ggctgcttgt acgtcttttg 3480
agaagtgtct cttcatgcct ttgtggcact ttaatgggat tattttttgc ttttttagtt 3540
aagtctctta tagattctgg atattagact tcttattgga tgcatagtgt gtgaatactc 3600
tcttcaccac tgtaggttgt ctgtttactc tattgatggc ttcttttgct gtgccgaagc 3660
atcttagttt aattagaaac cacctgccaa tttttgtttt tgttgcaatt gcttttgggg 3720
acttagtcat aaactctttg ccaaggtctg ggtcaagaag agtatttctt aggttttctt 3780
ctagaatttt gaaaggtctga atgtaaacat ttgcattttt aatgcattct gaggtagttt 3840
ttgtatatgt gaaaggtcta ctctcatttt ctttccctct ttctttcttt ctttcttttc 3900
ttctttcttt tctttctttc tttctttctt tctttctttc tttctttttg tctttctttc 3960
ttctttcttt tctctttctt tctctctttc tttttttttt ttgatggagt attgctctgt 4020
tgcccagggt gcagtgcagc ggcacgatct cggtcactg caacctctgc ctctcgggtt 4080
caactgattc tcttgcatca gccttccaag tagctgggat tataggcgcc cgccaccacg 4140
cccgactaat ttttgtattt ttagtagaga cgggggtgtg ccatgttggc caggctgggt 4200
tgaaactcct gacctcaaac gatctgcctg ccttggcctc ccaagtgtct gggattacag 4260
gtgtgagcca ctgtgccagc ccaagaatgt cattttctaa gaggtccaag aacctcaaga 4320
tattttggga ccttgagaag agaggaattc atacaggtat tacaagcaca gcctaattgc 4380
aaatcttttg catggcttgg ctccaagact ttatggctctt aaaagtctga tccaaaaatt 4440
tttataaaag ctccagctaa gctaccttaa aaggggcctg tatggctgat cactcttctt 4500
gctatacttt acacaataa acaggccaaa tataatgagg ccaaaattta ttttgcaaat 4560
aaattgttcc tgctatgatt tactcttggg aagaacaggg aaaatagaga aaattttaga 4620
ttgcacttga cttttttttc tgaattttta tatgtgccta caatttgagc taaattctga 4680
attattttct ggttgcaaaa actctctaaa gaagaacttg gtttctattg tcttcgtgac 4740
acatttatct ggctctttac tagaacagct ttcttggttt tgggtgtcta gcttggtgac 4800
cttacagttc tactcttcaa attattgtta tgtgtatctc atagttttcc ttcttttgag 4860
aaaactgaag ccatggtatt ctgaggacta gagatgactc aaacagagctg gtgaattctc 4920
tcataatgcaa tccactgggc tcgactctgt tcaaatgtct gatgcactgc tgctaaagct 4980
atcacattaa aacctcactc aaaggatcag ggaccatcat ggaagaggag gaaacatgaa 5040
attgtaagag ccagattcgg ggggtagagt gtggaggtca gagcaactcc acctgaaata 5100

FIGURE 6D

agaaggtaaa gcaacctatc ctgaaagcta acctgccatg gtgctcttg attaacctct 5160
 gttctaggaa gactgacagt ttgggtctgt gtcattgcc aaatctcatg ttaaatgtga 5220
 atccccagtg ttcggaagtg ggaacttggt gtgagtgatt cggtcatggg agtagatttt 5280
 cttctttgtg gtgttacagt gatagtgagt gattctctgt gagatctggt catttaaaag 5340
 tgtgtggccc ctcccctccc tctcttggtc ctccactgc catgtaagat acctgctcct 5400
 gctttgctt ctaccataag taaaagcccc ctgaggcctc ccagaagca gatgccacca 5460
 tgettcctgt acagcctgca gaacctcag ccaattaaac ctctttctg tataaattac 5520
 cagtcttgag tatctcttta cagcagtggt agaacggact aatacaaggg tctccaaat 5580
 tccaagtta tgtattcttt ctgccaat agcaggtatt taccataaat cctgtcctta 5640
 ggtcaaaaca ccttgatggc atcgtacttc aattgtctta cacattcctt ctgaatgact 5700
 cctcccctat ggcataaag ccttgggtct tgggggataa tggcagaggg gtccaccata 5760
 ttgtctggct gccacctgag acacggacat ggcttctgtt ggtaagtctc tattaatgt 5820
 ttctttctaa gaaactggat ttgtcagctt gttctttgg cctctcagct tctcagact 5880
 ttggggtagg ttgcacaacc ctgccacca cgaaacaaat gtttaatatg ataatatgg 5940
 atagatataa tccacataaa taaaagctct tggaggcccc tcaataattg ttaagagtg 6000
 aatgtgtcc aaagatggaa aatgtttgag aactactgtc ccagagattt tcttgagttc 6060
 tagagtgtg gaatatagaa cctggagctt ggtctctca gcttagaate agtagtatg 6120
 ggctgaagtc tgaagcttgg ctccagcagt ttggggttgg ctcccgagc acatatttga 6180
 catgttgca ctgtgatttg ggggttggtt ttgtctctga atctaatgt ctgtccttga 6240
 ggcatctaga atctgaaatc tgtggtcaga attctattat ctgagtagg acatctccag 6300
 tctgtgtct gccttctagg gctggagtct tagtcagtg acccggtctg gcatttcaac 6360
 ttcatataca ttgggctatc ttttggcca tgtttcaacc aaacaaccga ataaaccatt 6420
 agaaccttcc ccaacttccc tagtgcaat gttaaacctt ggatttctgt ttaatagtt 6480
 catatgaata atttcagct gatccaaatt tacattcctt ctaccgttat tctacacca 6540
 ccttaaaaat gcattcccaa tatattccct ggattctacc tatatatggt aatcctggt 6600
 ttgccagttt ctagtgcatt aacatactg atttaccct tttacttta aagtggaaat 6660
 aagatccct ctgcagagtt caggagttct caagatggcc ctacttctg acatcaattg 6720
 agatttcaag ggagtcgcca agatcatcct caggttcagt gattgtcgtt agccctcata 6780

FIGURE 6E

taactcaatg aaagctgtta tgctcatggc tatggtttat tacagcaaaa gaatagagat 6840
 gaaaatcttag caagggaaga gttgcatggg gcaaagacaa ggagagctcc aagtgcaag 6900
 attcctgttg tttctccca gtggtgcat ggaagcagt atcttctcca tacaatgatg 6960
 tgtgataata ttcagtgtat tgccaatcag ggaactcaac tgagccttga ttatattgga 7020
 gcttggttgc acagacatgt cgaccacctt catggctgaa ctttagtact tagccctcc 7080
 agcgtcttac agctgatagg ctgtaacca acattgtcac cataaatcac attgttagac 7140
 tatccagtgt ggcccaagct cccgtgtaaa cacaggcact ctaaacaggc aggatatttc 7200
 aaaaagcttag agatgacctc ccaggagctg aatgcaaaga cctggcctct ttgggcaag 7260
 agaactcttt accgcacct ctcttcaca gggttattgt gaggatcaaa tgtggtcatg 7320
 tgtgtgagac accagcacat gtctggctgt ggagagtac ttctatgtgt gctaaccattg 7380
 ctgagtgccta agaaagtatt aggcattgct ttcagcactc acagatgtct atctaactct 7440
 cacaacatgg ctacagggtg ggccactact gcctcatttg acagaggaaa ggactgtgga 7500
 taagaagggg gtgaccaata ggctcagatc attctggatg caaggggctc cagaggacca 7560
 tgattagaca ttgtctgcag agaaattatg gctggatgtc tctgccccgg aaagggggat 7620
 gcactttcct tgacccccta tctcagatct tgactttgag gttatctcag acttctccta 7680
 tgataccagg agcccatcat aatctctctg tgtcctctcc cttctctcag tcttactgcc 7740
 cactcttccc agctccatct ccagctggcc aggtgtagcc acagtaccta actctttgca 7800
 gagaactata aatgtgtatc ctacagggga gaaaaaaaa aagaactctg aaagagctga 7860
 cattttaccg acttgcaaac acataagcta atctgccagt ttgtgctgg tagaactcat 7920
 gagactcctg ggtcagaggc aaaagatttt attaccaca gctaaggagg cagcatgaac 7980
 tttgtgtcca catttgttca ctttgcccc caattcatat gggatgatca gagcagtcca 8040
 ggtggatgga cacagggggt ttgtggcaag gtgagcaacc taggcttaga aatcctcaat 8100
 cttataagaa ggtactagca aacttgtcca gtctttgtat ctgacggaga tattatcttt 8160
 ataattgggt tgaagcaga cctactctgg aggaacatat tgtatttatt gtcttgaaca 8220
 gtaaacaaat ctgctgtaaa atagacgtta actttattat ctaaggcagt aagcaaacct 8280
 agatctgaag gcataccat cttgcaagge tatctgctgt acaaatatgc ttgaaaagat 8340
 ggtccagaaa agaaaacggt attattgect ttgtcagaa gacacacaga aacataagag 8400
 aacctggaa aattgtctcc caacctgtt caccagagc cttccactct tgtctgcagg 8460
 acagtcttaa catccatca ttagtgtgtc taccacatct ggcttaccg tgcctaacca 8520

FIGURE 6F

agatttctag gtccagtgtc ccaccatgtt tggcagtgcc ccactgccaa cccagaata 8580
 agggagtgct cagaattccg aggggacatg ggtggggate agaacttctg ggcttgagtg 8640
 cagagggggc ccatactcct tggttccgaa ggaggaagag gctggagggtg aatgtccttg 8700
 gaggggagga atgtgggttc tgaactctta aatccccaag ggaggaagact ggtaagggtcc 8760
 cagcttccga ggtactgacg tgggaatggc ctgagagggtc taagaatccc gtatcctcgg 8820
 gaaggagggg ctgaaattgt gaggggttga gttgcagggtg tttgttagct tgagactcct 8880
 tgggtgggtcc ctgggaagca aggaactggaa ccattggctc cagggtttgg tgtgaaggta 8940
 atgggacttc ctgattctca aagggtcaga ggactgagag ttgcccatgc tttgatcttt 9000
 ccactactc cttactccac ttgagggtaa tcactactc ttctagtcc acaagagtgc 9060
 gctgcgcga gtataatctg cacatgtgcc atgtcccgag gctggggcca tcattccactc 9120
 atcattcagc atctgcgcga tgcggggcag gcccggccca tgacgtcatg tagctgcgac 9180
 tatccctgca gcgcgcctct ccctgcacgt cccaaccatg gagctgtgga cgtgcgtccc 9240
 ctggtggatg tggcctgcgt ggtgccaggc cggggcctgg tgtccgataa agatcctaga 9300
 accacagga accaggactg aaagggtgta gagaatggcc atatgtcgct gtccatgaaa 9360
 tctcaaggac ttctgggtg agggcacagg agcctgaact tacgggtttg cccagtgcca 9420
 ctgtctctcc aagtgaagt cccagatacg aggcactgtg ccagcatcag cttcatctgt 9480
 accacatctt gtaacaggga ctaccaggga ccctgatgaa caccatgggtg tgtgcaggaa 9540
 gaggggggtg aggcattggc tcctgtgtgg tcagagccca gagggggcca tgacgggtgg 9600
 ggaggaggct gtggactggc tcgagaagtg ggaatgtggt gtgtttgatt tcctttggcc 9660
 agataaagt ctggatatag cattgaaaac ggagtatgaa gaccagttag aatggagggt 9720
 caggttggag ttgagttaca gatggggtaa aattctgctt cggatgagtt tggggattgg 9780
 caatctaaag gtggtttggg atggcatggc ttgggatgg aaatagggtt gtttttatgt 9840
 tggctgggaa ggggtgtggg attgaattgg gaatgaagta ggtttagttt tggagataga 9900
 atacatggag ctggctattg catgcgagga tgtgcattag tttggtttga tctttaata 9960
 aaggaggcta ttagggttgt cttgaattag attaagttgt gttgggttga tgggtgtggc 10020
 ttgtgggtga tgtggttga ttgggctgtg ttaaattggt ttgggtcagg ttttggttga 10080
 ggttatcatg gggatgagga tatgcttggg acatggatcc aggtgttct cattcaagct 10140
 gaggcaaatt tcctttcaga cgggtcattcc agggaaacgag tggttgtgtg ggggaaatca 10200

FIGURE 6G

ggccactggc tgtgaatc cctctatcct ggtcttgaat tgtgattatc tatgtccatt 10260
 ctgtctcctt cactgtactt ggaattgacg tggctattca gctggaaatg ggggaagatt 10320
 ttgtcaaatt cttgagacac agctgggtct ggtacagcgt aagccttctt cctgttttta 10380
 ttgaacagat gaaatcacat tttttttttt aaaatcacag aaatcttata gagttaacag 10440
 tggactctta taataagagt taacaccagg actcttattc ttgattcttt cctgagacac 10500
 caaaatgaga tttctcaatg ccaccttaat tctttttttt tttttttttt tttttgagac 10560
 acagtctggg tcttttcttc tgtcactcag gctggagcgc agtgggtgtg tcatagctca 10620
 ctgaaccctt gacctctctg acttaaggga tctctctgct tcagcctcct gagtagatgg 10680
 ggctacaggg gcttgccacc acacctggct aattaaattt tttttttttt ttttagaga 10740
 aagggtctca ctttgttgcc ctggtctgac ttgaactctt gaettcaagt gattcttcag 10800
 ccttggaetc ccaaagcact gggattgctg gcatgagcca ctacccgtgc ctggcttgca 10860
 gcttaactctt ggaagtata aacctggctc ctgatatcta gacatttcag tgagaaggag 10920
 gcatctgatt ttgcatgagg acaattctga cctaggaggg caggtcaaca ggaatccccg 10980
 ctgtacctgt acgttgtaca ggcattgaga atgaggagtg aggagggcgt accggaacc 11040
 catattgttt agtggacatt ggattttgaa ataataaggga acttggtctg ggagagtcac 11100
 atttctggat tggacaatat gtggtatcac aaggttttat gatgaggag aaatgtatgt 11160
 ggggaacat tttctgagtg tgggaagtga agaatacagag agtagctgaa tgccaacgct 11220
 tctatttcag gaacatggta agttggaggc ccagctctcg ggctcagacg ggtataggga 11280
 ccagggaatc tcacaatccg atcattctga tatttcaggg catattaggt ttggggtgca 11340
 aagggaagtac ttgggactta ggcacatgag actttgtatt gaaaatcaat gattggggct 11400
 ggccgtgtg ctcacgcctg taatctctac actttgggag accgaagtg gaggatggct 11460
 tgatctcaag agttggacac cagcctaggc aacatggcca gacctctct ctacaaaaaa 11520
 attaaaaatt agctggatgt ggtgtgtcat gcttgggtc tcagctatcc tggaggctga 11580
 gacaggagaa tcggttgagt ctgggagttc aagcctacag ggagctgcga tcacgcctcg 11640
 gcactccagc ctgggaacaa gagtgagact gtctcagaat ttttttaaaa aagaatcagt 11700
 gatcatccca acccctgttg ctgttcacac tgagcctgcc ttctctggct ttgttcccta 11760
 gatcacatct ccatgatcca taggcctcgc ccaatctgac ctacacacct gggaaatgct 11820
 ccagactgat ctagtatgtg tggaacagca agtgcctggc ctccctcccc ttccacagct 11880
 ctgggtgtg gagggggttg tccagcctcc agcagcatgg ggagggcctt ggtcagcacc 11940

FIGURE 6H

taggtgccaa cagggcaagg gcggggcct ggagaatgaa ggctttatag ggctctcag 12000
ggaggccccc cagccccaaa ctgcaccacc tggccgtgga caccggt 12047

FIGURE 7

```

cgagcggccc ctcagcttcg gcgccagcc cgcgaagct cccgtgacc actagagggc 60
gggaggagct cctggccagt ggtggagagt ggcaaggaag gaccctaggg ttcacggag 120
cccaggttta ctccttaag tggaaatttc ttccccact cctccttggc tttctccaag 180
gaggggaaccc aggtcgtcg aaagtcggcg tgggcgggg actgtgggtt cagggggagaa 240
cggggtgtgg aacgggacag ggagcgggta gaagggtggg gctattccgg gaagtgggtg 300
ggggagggag cccaaaacta gcacctagtc cactcattat ccagccctct tattctcgg 360
ccgctctgct tcagtggacc cggggagggc ggggaagtgg agtgggagac ctaggggtgg 420
gcttcccgac ctgctgtac aggacctga cctagctggc ttgttcccc atccccagt 480
tagttgttgc cctgaggcta aaactagagc ccaggggccc caagttccag actgccctc 540
ccccctccc cggagccagg gagtgggttg tgaaggggg aggcagctg gagaacaaac 600
gggtagtacg ggggttgagc gattagagcc ctgtaccct acccaggaat ggttggggag 660
gaggaggaag aggtaggagg taggggaggg ggcggggtt tgtcaactgt cacctgctcg 720
ctgtcctcag ggcgggaggg cggggagtgg ggggaccggt ataaagcggg aggcgcctgt 780
gcccgcteca cctctcaagc agccagcgcc tgccgaatc tgtctgccc cctccccacc 840
catttcacca ccaccatg
858

```

FIGURE 8

aagcttccac aagtcattt agcctctcca gtattgctga tgaatccaca gttcaggttc 60
aatggcggtc aaaacttgat caaaaatgac cagactttat attcttacac caacatctat 120
ctgattggag gaatggataa tagtcatcat gtttaacat ctaccattcc agttaagaaa 180
atatgatagc atctgttct tagtcttttt cttaataggg acataaagcc cacaataaaa 240
aatatgcctg aagaatggga caggcattgg gcattgtcca tgctagtaa agtactccaa 300
gaacctattt gtatactaga tgacacaatg tcaatgtctg tgtacaactg ccaactggga 360
tgcaagacac tgcccatgcc aatcatcctg aaaagcagct ataaaaagca ggaagctact 420
ctgcaccttg tcagtgggt ccagatacct acag 454

FIGURE 9

```

g atg acc ggc tca acc atc gcg ccc aca acg gac tat cgc aac acc act 49
Met Thr Gly Ser Thr Ile Ala Pro Thr Thr Asp Tyr Arg Asn Thr Thr
1 5 10 15

gct acc gga cta aca tct gcc cta aat tta ccc caa gtt cat gcc ttt 97
Ala Thr Gly Leu Thr Ser Ala Leu Asn Leu Pro Gln Val His Ala Phe
20 25 30

gtc aat gac tgg gcg agc ttg gac atg tgg tgg ttt tcc ata gcg ctt 145
Val Asn Asp Trp Ala Ser Leu Asp Met Trp Trp Phe Ser Ile Ala Leu
35 40 45

atg ttt gtt tgc ctt att att atg tgg ctt att tgt tgc cta aag cgc 193
Met Phe Val Cys Leu Ile Ile Met Trp Leu Ile Cys Cys Leu Lys Arg
50 55 60

aga cgc gcc aga ccc ccc atc tat agg cct atc att gtg ctc aac cca 241
Arg Arg Ala Arg Pro Pro Ile Tyr Arg Pro Ile Ile Val Leu Asn Pro
65 70 75 80

cac aat gaa aaa att cat aga ttg gac ggt ctg aaa cca tgt tct ctt 289
His Asn Glu Lys Ile His Arg Leu Asp Gly Leu Lys Pro Cys Ser Leu
85 90 95

ctt tta cag tat gat taa 307
Leu Leu Gln Tyr Asp
100

```

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/US 99/20718

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/86 C12N5/10 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 698 443 A (HENDERSON DANIEL ROBERT ET AL) 16 December 1997 (1997-12-16) cited in the application the whole document ---	1-17, 23-28
Y	SEMENZA G. L. ET AL.: "Hypoxia response elements in the aldolase A, enolase 1, and lactate dehydrogenase A gene promoters contain essential binding sites for hypoxia-inducible factor 1." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, no. 51, 1996, pages 32529-32537, XP002129236 the whole document, especially Fig. 10 --- -/-	1-10, 14-17, 23-28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document published on or after the international filing date
- *L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (see specification)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

31 January 2000

Date of mailing of the international search report

11/02/2000

Name and mailing address of the ISA
European Patent Office, P.B. 5816 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mandl, B

INTERNATIONAL SEARCH REPORT

Inte onal Application No
PCT/US 99/20718

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 13508 A (DANA FARBER CANCER INST INC ;KAELIN WILLIAM JR (US); FINE HOWARD A) 2 April 1998 (1998-04-02) page 3, last paragraph - paragraph 2 page 9, line 1 - line 11 ---	1-8,11, 12, 14-17, 23-28
Y	WO 98 06864 A (US HEALTH ;MOONEN CHRIT (US)) 19 February 1998 (1998-02-19) page 10, line 36 -page 12, line 8 ---	1-8, 13-17, 23-28
A	EP 0 845 537 A (CHIRON CORP) 3 June 1998 (1998-06-03) page 22, line 31 - line 34 ---	18-22
A	WO 96 17053 A (GENETIC THERAPY INC ;HALLENBECK PAUL L (US); CHANG YUNG NIEN (US);) 6 June 1996 (1996-06-06) cited in the application the whole document ---	1-28
A	RINSCH C. ET AL.: "A GENE THERAPY APPROACH TO REGULATED DELIVERY OF ERYTHROPOIETIN AS A FUNCTION OF OXYGEN TENSION." HUMAN GENE THERAPY, vol. 8, no. 16, November 1997 (1997-11), pages 1881-1889, XP000867701 the whole document ---	9,10
A	ZWICKER J. ET AL.: "CELL CYCLE REGULATION OF THE CYCLIN A, CDC25C AND CDC2 GENES IS BASED ON A COMMON MECHANISM OF TRANSCRIPTIONAL REPRESSION" EMBO JOURNAL, vol. 14, no. 18, 1 January 1995 (1995-01-01), pages 4514-4522, XP002038970 ISSN: 0261-4189 the whole document -----	11,12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/20718

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 27 and 28, as far as in vivo application is concerned, are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

Info onl Application No
PCT/US 99/20718

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5698443 A	16-12-1997	AU 6393296 A	30-01-1997
		CA 2222457 A	16-01-1997
		EP 0844888 A	03-06-1998
		JP 11508770 T	03-08-1999
		WO 9701358 A	16-01-1997
		US 5871726 A	16-02-1999
WO 9813508 A	02-04-1998	AU 4592697 A	17-04-1998
WO 9806864 A	19-02-1998	AU 4241697 A	06-03-1998
		EP 0922110 A	16-06-1999
EP 0845537 A	03-06-1998	AU 5915394 A	16-06-1994
		AU 648261 B	21-04-1994
		AU 6185390 A	03-04-1991
		AU 694685 B	23-07-1998
		AU 7058496 A	16-01-1997
		CA 2066053 A	19-02-1991
		EP 0487587 A	03-06-1992
		JP 4507196 T	17-12-1992
		US 5662896 A	02-09-1997
		WO 9102805 A	07-03-1991
		US 5716826 A	10-02-1998
		US 5591624 A	07-01-1997
		US 5691177 A	25-11-1997
		US 5716832 A	10-02-1998
		US 5888502 A	30-03-1999
		US 5997859 A	07-12-1999
		US 5856185 A	05-01-1999
		US 5716613 A	10-02-1998
		US 5851529 A	22-12-1998
		US 5830458 A	03-11-1998
WO 9617053 A	06-06-1996	AU 4504396 A	19-06-1996
		CA 2206179 A	06-06-1996
		EP 0791050 A	27-08-1997
		FI 972225 A	16-07-1997
		JP 10509880 T	29-09-1998
		NO 972373 A	25-07-1997
		US 5998205 A	07-12-1999